

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>4</sup> :</b> C07D 215/56, 401/04, 471/04 C07D 451/04, A61K 31/44 A61K 31/47, C07C 51/00 // (C07D 471:04, 221:00, 221:00)	<b>A2</b>	<b>(11) International Publication Number:</b> WO 89/ 06649		
		<b>(43) International Publication Date:</b> 27 July 1989 (27.07.89)		
<table border="0" style="width: 100%;"><tr><td style="width: 50%; vertical-align: top;"><b>(21) International Application Number:</b> PCT/US89/00278 <b>(22) International Filing Date:</b> 23 January 1989 (23.01.89) <b>(31) Priority Application Numbers:</b> 147,462 280,924 <b>(32) Priority Dates:</b> 25 January 1988 (25.01.88) 9 December 1988 (09.12.88) <b>(33) Priority Country:</b> US <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> US 280,924 (CIP) Filed on 9 December 1988 (09.12.88) <b>(71) Applicant (for all designated States except US):</b> WARN- ER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).</td><td style="width: 50%; vertical-align: top;"><b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DOMAGALA, John, Michael [US/US]; 776 Georgetown, Canton, MI 48188 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton, MI 48187 (US). KIELY, John, Steven [US/US]; 4138 Sunset Court, Ann Arbor, MI 48103 (US). <b>(74) Agents:</b> ANDERSON, Elizabeth, M.; Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, MI 48105 (US) et al. <b>(81) Designated States:</b> AT, AT (European patent), AU, BE (European patent), CH (European patent), DE, DE (European patent), DK, FI, FR (European patent), GB, GB (European patent), IT (European patent), JP, KR, LU, LU (European patent), NL, NL (European patent), NO, SE, SE (European patent), US. <b>Published</b> <i>Without international search report and to be repu- blished upon receipt of that report.</i></td></tr></table>			<b>(21) International Application Number:</b> PCT/US89/00278 <b>(22) International Filing Date:</b> 23 January 1989 (23.01.89) <b>(31) Priority Application Numbers:</b> 147,462 280,924 <b>(32) Priority Dates:</b> 25 January 1988 (25.01.88) 9 December 1988 (09.12.88) <b>(33) Priority Country:</b> US <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> US 280,924 (CIP) Filed on 9 December 1988 (09.12.88) <b>(71) Applicant (for all designated States except US):</b> WARN- ER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).	<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DOMAGALA, John, Michael [US/US]; 776 Georgetown, Canton, MI 48188 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton, MI 48187 (US). KIELY, John, Steven [US/US]; 4138 Sunset Court, Ann Arbor, MI 48103 (US). <b>(74) Agents:</b> ANDERSON, Elizabeth, M.; Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, MI 48105 (US) et al. <b>(81) Designated States:</b> AT, AT (European patent), AU, BE (European patent), CH (European patent), DE, DE (European patent), DK, FI, FR (European patent), GB, GB (European patent), IT (European patent), JP, KR, LU, LU (European patent), NL, NL (European patent), NO, SE, SE (European patent), US. <b>Published</b> <i>Without international search report and to be repu- blished upon receipt of that report.</i>
<b>(21) International Application Number:</b> PCT/US89/00278 <b>(22) International Filing Date:</b> 23 January 1989 (23.01.89) <b>(31) Priority Application Numbers:</b> 147,462 280,924 <b>(32) Priority Dates:</b> 25 January 1988 (25.01.88) 9 December 1988 (09.12.88) <b>(33) Priority Country:</b> US <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> US 280,924 (CIP) Filed on 9 December 1988 (09.12.88) <b>(71) Applicant (for all designated States except US):</b> WARN- ER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).	<b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> DOMAGALA, John, Michael [US/US]; 776 Georgetown, Canton, MI 48188 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton, MI 48187 (US). KIELY, John, Steven [US/US]; 4138 Sunset Court, Ann Arbor, MI 48103 (US). <b>(74) Agents:</b> ANDERSON, Elizabeth, M.; Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, MI 48105 (US) et al. <b>(81) Designated States:</b> AT, AT (European patent), AU, BE (European patent), CH (European patent), DE, DE (European patent), DK, FI, FR (European patent), GB, GB (European patent), IT (European patent), JP, KR, LU, LU (European patent), NL, NL (European patent), NO, SE, SE (European patent), US. <b>Published</b> <i>Without international search report and to be repu- blished upon receipt of that report.</i>			
<b>(54) Title:</b> ANTIBACTERIAL AGENTS				
<b>(57) Abstract</b>				
<p>Novel naphthyridine-, and quinolinecarboxylic acids as antibacterial agents are described as well as methods for their manufacture, formulation, and use in treating bacterial infections including the description of certain novel intermediates used in the manufacture of the antibacterial agents.</p>				

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				


## ANTIBACTERIAL AGENTS

## CROSS-REFERENCE TO RELATED APPLICATIONS

This is a continuation-in-part of United States Application Serial No. 147,462 filed January 25, 1988.


## BACKGROUND OF THE INVENTION

US Patent 4,341,784 discloses certain substituted 7-(3-amino-1-pyrrolidinyl)-1-ethyl-6-fluoro-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acids  
5 having the general formula:



The compounds are disclosed to have antibacterial activity.

The Journal of Medicinal Chemistry, 23, 1358 (1980) discloses certain substituted quinoline-3-carboxylic acids having the structural formula  
10



wherein -N $\bigcirc$  may be pyrrolidinyl. See also US


Patent 4,146,719. The compounds are disclosed to have antibacterial activity.

Certain 7-heterocyclic substituted 1,8-naphthyridines are disclosed in Eur. J. Med. Chem. - Chemica  
15

-2-


Therapeutica, 29, 27 (1977). US Patents 3,753,993 and 3,907,808 disclose certain 7-pyridylquinolones.

European Patent Applications 229,635 and 206,101 cover certain 1,8-bridged-1,4-dihydro-4-quinolinones having the formula




wherein  $X_1$  is hydrogen,  $\text{NO}_2$ , 1-3C alkyl or halogen;  $X_2$  is halogen, 1-3C-alkyl, 1-3C-alkylsulphenyl or optionally substituted phenylsulphenyl;  $X_5$  is hydrogen, halogen or methyl.

10 US Patent 4,774,246 discloses certain substituted 1-phenyl-6,8-difluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-quinoline-3-carboxylic acids of general formula




15 US Patent 4,704,459 discloses a process for certain 1-substituted aryl-1,4-dihydro-4-oxonaphthyridine derivatives of general formula

-3-



US Patent 4,649,144 discloses certain  
1,8-naphthyridine derivatives of general formula




US Patent 4,571,396 discloses certain  
naphthyridine-quinoline-, and benzoxazine-carboxylic  
5 acids with a bridged side-chain at the seven-position.  
Copinging US application 080,113 discloses  
certain naphthyridine-, quinoline-, and benzoxazine-  
carboxylic acids with a bridged side-chain at the  
seven-position and a hydrogen, fluoro or amino at the  
10 five-position.

The references teach that these compounds possess  
antibacterial activity.

#### SUMMARY OF THE INVENTION


One aspect of the present invention is a compound  
15 of Formula I

-4-









or a pharmaceutically acceptable acid addition or base salt thereof wherein

Z is



or



wherein  $R_4$  is hydrogen, alkyl of from one to  
 5 four carbon atoms, or cycloalkyl of from three to

-5-

six carbon atoms, R' is hydrogen, hydroxyl, alkyl of from one to four carbon atoms, phenyl or phenyl substituted by halogen, alkyl or alkoxy, n is an integer of from 0 to 4, R<sub>5</sub> and R<sub>6</sub> are each  
5 independently hydrogen, lower alkyl or cycloalkyl; X is CH, CF, CCl, CBr, N, CCF<sub>3</sub>, CNH<sub>2</sub>, CNO<sub>2</sub>, CR, or COR' wherein R is lower alkyl and R'' is hydrogen or lower alkyl;

10 R<sub>3</sub> is lower straight, branched, or cyclic alkyl of from one to three carbon atoms;

R<sub>2</sub> is alkyl of from one to four carbon atoms, vinyl, haloalkyl, hydroxyalkyl of from two to four carbon atoms, cycloalkyl of from three to six carbon atoms, phenyl or phenyl substituted by  
15 halogen, alkyl, NH<sub>2</sub> or OH;

R<sub>1</sub> is hydrogen, alkyl of from one to six carbon atoms, or a cation.

The preferred compounds of this invention are those wherein X is CH, CF, CCl, or N.


20 Also preferred compounds of the invention are those wherein R<sub>2</sub> is cyclopropyl, ethyl, or 2,4-difluorophenyl.

Other preferred compounds of the invention are those wherein R<sub>3</sub> is methyl, ethyl, isopropyl, or  
25 cyclopropyl.

Other preferred compounds of this invention are those wherein R<sub>1</sub> is hydrogen or a pharmaceutically acceptable base salt such as a metal or amine salt.

Other preferred compounds of this invention are  
30 those wherein Z is

-6-














wherein  $R_4$  is hydrogen or methyl,  $R'$  is hydrogen or methyl,  $n$  is 0, 1, or 2,  $R_5$  and  $R_6$  are each independently hydrogen or methyl.

Particularly preferred compounds are those where  
 5 Z is selected from the group consisting of



-7-




or



-8-

Most preferred compounds include those wherein X is CH, CF, CCl; R<sub>2</sub> is cyclopropyl; R<sub>3</sub> is CH<sub>3</sub>, Et; R<sub>1</sub> is H; and Z is



Particularly preferred compounds of the invention are compounds having the names:

- 5 1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,
- 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-
- 10 carboxylic acid,
- 1-cyclopropyl-7-[3-(ethylamino)methyl]-1-pyrrolidinyl]-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-
- 15 cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-7-[3-methyl-1-piperazinyl]-4-oxo-3-quinolinecarboxylic acid,
- 20 1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,
- 1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,
- 25 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinoline-carboxylic acid,

5 7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinoline-carboxylic acid,

7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

10 7-[3-(endo-amino)-8-azabicyclo[3.2.1]oct-8-yl]-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

and pharmaceutically acceptable acid addition or base salts thereof.

15 Other preferred compounds of the invention are compounds having the names:

1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-ethyl-7-[3-methyl-1-piperazinyl]-4-oxo-3-quinolinecarboxylic acid,

20 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-5-ethyl-6,8-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid,

7-(3-amino-1-pyrrolidinyl)-1,5-dicyclopropyl-6,8-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid,

25 1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

30 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

7-(3-amino-1-pyrrolidinyl)-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

35 8-chloro-1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

-10-

8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,

5 8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

10 7-(3-amino-1-pyrrolidinyl)-8-bromo-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

8-bromo-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-8-bromo-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

20 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-8-hydroxy-5-methyl-4-oxo-3-quinolinecarboxylic acid,

1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-8-hydroxy-5-methyl-4-oxo-3-quinolinecarboxylic acid,

25 1-cyclopropyl-6-fluoro-1,4-dihydro-8-hydroxy-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,

7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-5-methyl-4-oxo-3-quinolinecarboxylic acid,

30 1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-8-methoxy-5-methyl-4-oxo-3-quinolinecarboxylic acid,

35 1-cyclopropyl-6-fluoro-1,4-dihydro-8-methoxy-7-(3-methyl-1-piperazinyl)-5-methyl-4-oxo-3-quinolinecarboxylic acid,

-11-

7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-8-nitro-4-oxo-3-quinolinecarboxylic acid,

5 1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-8-nitro-4-oxo-3-quinolinecarboxylic acid,

7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-8-nitro-4-oxo-3-quinolinecarboxylic acid,

10 8-amino-7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

8-amino-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

15 8-amino-7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

20 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-5-methyl-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid,

1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-1,8-naphthyridine-3-carboxylic acid,

25 1-cyclopropyl-5-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-1,8-naphthyridine-3-carboxylic acid.

7-(3-amino-1-pyrrolidinyl)-6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

30 6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

35

-12-

- 7-(3-amino-1-pyrrolidinyl)-1-ethyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
1-ethyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-  
5 carboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-6,8-difluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
6,8-difluoro-1-(2-fluoroethyl)-1,4-dihydro-5-  
10 methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-1-vinyl-3-quinolinecarboxylic acid,  
15 6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid,  
6,8-difluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-1-vinyl-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-  
20 difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
25 7-(3-amino-1-pyrrolidinyl)-1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-  
30 fluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
6-fluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

- 7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 5 7-(3-amino-1-pyrrolidinyl)-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1-vinyl-3-quinolinecarboxylic acid,
- 6-fluoro-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-1-vinyl-3-quinolinecarboxylic acid,
- 8-chloro-7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 10 8-chloro-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,
- 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-8-chloro-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 15 8-chloro-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,
- 20 8-chloro-1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-piperazinyl-3-quinolinecarboxylic acid,
- 7-(3-amino-1-pyrrolidinyl)-8-chloro-1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 25 8-chloro-1-ethyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 7-(3-amino-1-pyrrolidinyl)-8-chloro-6-fluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 30 8-chloro-6-fluoro-1-(2-fluoroethyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,
- 7-(3-amino-1-pyrrolidinyl)-8-chloro-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1-vinyl-3-quinolinecarboxylic acid,
- 35

-14-

- 8-chloro-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
5 1-cyclopropyl-6-fluoro-1,4-dihydro-5,8-dimethyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
1-cyclopropyl-6-fluoro-1,4-dihydro-5,8-dimethyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
10 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
15 7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
20 6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5,8-dimethyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
25 7-(3-amino-1-pyrrolidinyl)-1-ethyl-6-fluoro-1,4-dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid,  
1-ethyl-6-fluoro-1,4-dihydro-5,8-dimethyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
30 7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
1-cyclopropyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
35



-15-

1-cyclopropyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

5 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

10 7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

6-fluoro-1-(2,4-difluorophenyl)-8-trifluoromethyl-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

15 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

7-(3-amino-1-pyrrolidinyl)-1-ethyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

20 1-ethyl-6-fluoro-8-trifluoromethyl-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

25 1-ethyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,

6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,

30 6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,

6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid, and

35

-16-


7-[3-(ethylamino)methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid.

The invention further includes certain novel  
5 intermediate compounds having the names  
1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid ethyl ester,  
2,3,4,5-tetrafluoro-6-methylbenzoic acid,  
2,3,4,5-tetrafluoro-6-methylbenzoyl chloride,  
10 ethyl 3-(2,3,4,5-tetrafluoro-6-methylphenyl)- $\beta$ -oxo-propanoate,  
ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-ethoxyacrylate,  
2-(2,4,5-trifluoro-3-trimethylsilylphenyl)-4,4-  
15 dimethyl-2-oxazoline,  
ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-cyclopropylaminoacrylate,  
2-(2,4,5-trifluoro-6-methyl-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline,  
20 2-(2,4,5-trifluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline,  
2-(3-chloro-2,4,5-trifluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline,  
2-(3-bromo-2,4,5-trifluoro-6-methylphenyl)-4,4-  
25 dimethyl-2-oxazoline,  
2-(2,4,5-trifluoro-3-hydroxy-6-methylphenyl)-4,4-dimethyl-2-oxazoline,  
2-(2,4,5-trifluoro-6-methyl-3-nitrophenyl)-4,4-dimethyl-2-oxazoline,  
30 2-(2,4,5-trifluoro-3,6-dimethylphenyl)-4,4-dimethyl-2-oxazoline,  
2-[2,4,5-trifluoro-3-(trifluoromethyl)-6-methylphenyl)-4,4-dimethyl-2-oxazoline,  
2,6-dichloro-5-fluoro-4-methyl-3-pyridinecarboxylic  
35 acid,


-17-

3-chloro-2,4,5-trifluoro-6-methylbenzoic acid,  
3-bromo-2,4,5-trifluoro-6-methylbenzoic acid,  
2,4,5-trifluoro-6-methyl-3-nitrobenzoic acid, and  
2,4,5-trifluoro-3,6-dimethylbenzoic acid.

- 5 Another aspect of the instant invention is the following process for preparing compounds of Formula I




wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, Z, n, R', and X are as defined above which comprises reacting a compound of Formula II




- 10 with an amine corresponding to the group Z  
wherein all of the above terms are as defined above in  
Formula I and L is a leaving group which may  
preferably be fluorine or chlorine.

- Yet another aspect of the instant invention is a  
15 process for preparing compounds of formula

-18-




wherein R is alkyl which comprises reacting a penta-fluorooxazoline with alkyl lithium producing a compound of formula




followed by acidic hydrolysis.

- 5 Yet another aspect of the present invention is a process for preparing compounds of formula




wherein R is alkyl which comprises  
(a) reacting a compound of formula


-19-



with a base and trimethylsilyl chloride producing a compound of formula



(b) reacting that compound with a base and an alkyl halide producing a compound of formula




- 5           (c) removing the SiMe<sub>3</sub> and  
             (d) hydrolyzing the resulting compound.


Yet another aspect of the present invention is a process for preparing naphthyridines of Formula I by

(a) reacting a compound of formula


-20-




with oxalyl chloride and dimethylformamide and quenching with alcohol to produce the corresponding ester



(b) reducing the double bond to produce a  
5 compound of formula



(c) treating the compound from step (b) with a base, then methyl iodide to produce the alkylated compound



-21-

(d) reintroducing the double bond and reacting the resulting naphthyridine with the desired amine by known means.

5 The invention also includes a pharmaceutical composition which comprises an antibacterially effective amount of a compound having structural Formula I and the pharmaceutically acceptable salts thereof in combination with a pharmaceutically acceptable carrier.

10 The invention further includes a method for treating bacterial infections in a mammal which comprises administering an antibacterially effective amount of the above defined pharmaceutical composition to a mammal in need thereof.

15 DETAILED DESCRIPTION

The compound of the invention having the structural Formula I may be readily prepared by treating a corresponding compound having the Formula II above with the desired cyclic amine as defined by Z. For purposes of this reaction, the alkylamine substituent of Z may, if desired, be protected by a group which renders it substantially inert to the reaction conditions. Thus, for example, protecting groups such as the following may be utilized:

25 carboxylic acyl groups such as formyl, acetyl, trifluoroacetyl;  
alkoxycarbonyl groups such as ethoxycarbonyl, t-butoxycarbonyl,  $\beta,\beta,\beta$ -trichloroethoxycarbonyl,  
30  $\beta$ -iodoethoxycarbonyl;  
aryloxycarbonyl groups such as benzyloxycarbonyl, p-methoxybenzyloxycarbonyl, phenoxycarbonyl;  
silyl groups such as trimethylsilyl; and groups such as trityl, tetrahydropyranyl, vinyloxycarbonyl,

-22-

o-nitrophenylsulfenyl, diphenylphosphinyl, p-toluenesulfonyl, and benzyl may all be utilized. The protecting group may be removed after the reaction between a compound as defined by Formula II and Z if  
5 desired, by procedures known to those skilled in the art. For example, the ethoxycarbonyl group may be removed by acid or base hydrolysis and the trityl group may be removed by hydrogenolysis.

The reaction between the compound Formula II and  
10 a suitably protected compound as defined by Z may be performed with or without a solvent, preferably at elevated temperature for a sufficient time so that the reaction is substantially complete. The reaction is preferably carried out in the presence of an acid  
15 acceptor such as an alkali metal or alkaline earth metal carbonate or bicarbonate, a tertiary amine such as triethylamine, pyridine, or picoline. Alternatively an excess of the compound of Formula VI may be utilized as the acid acceptor.

~~20 Convenient solvents for this reaction are~~  
nonreactive solvents such as acetonitrile, tetrahydrofuran, ethanol, chloroform, dimethylsulfoxide, dimethylformamide, pyridine, picoline, water, and the like. Solvent mixtures may also be utilized.

25 Convenient reaction temperatures are in the range of from about 20° to about 150°C; higher temperatures usually require shorter reaction times.


The removal of the protecting group may be accomplished either before or after isolating the  
30 product, I. Alternatively, the protecting group need not be removed.

The compounds of the invention of Z are either known compounds or they may be prepared from known starting materials by standard procedures or by




-23-

variations thereof. For example, 3-pyrrolidinemethanamines having the formula D



D

- 5 may be readily prepared from the known starting material methyl 5-oxo-1-(phenylmethyl)-3-pyrrolidine-carboxylate, A, [J. Org. Chem., 26, 1519 (1961)] by the following reaction sequence.



-24-


The compound wherein  $R_3$  is hydrogen, namely 3-pyrrolidinemethanamine, has been reported in J. Org. Chem., 26, 4955 (1961).

Thus Compound A may be converted to the  
5 corresponding amide B by treatment with  $R_3NH_2$ ; for example, a saturated solution of ethylamine in an alkanol such as methyl alcohol may be utilized. The diamide B may next be reduced to produce the corresponding diamine C. This reduction may be  
10 carried out using lithium aluminum hydride, for example, in a convenient solvent such as tetrahydrofuran. Compound C may next be debenzylated, for example using hydrogen and 20% palladium on carbon catalyst to produce the diamine D. Alternatively,  
15 when  $R = H$  in C, the primary amine function may be protected with a group  $R_4$  as defined, hereinabove. For example, the primary amine function may be acylated with an acyl halide such as acetyl chloride by well known procedures. The primary amine function  
20 of C may also be converted to a carbamate ester such as the ethyl ester by treatment with ethyl chloroformate in the presence of a strong base such as 1,8-diazabicyclo[5.4.0]undec-7-ene in a convenient solvent such as methylene chloride. The benzyl group  
25 may next be removed, for example as described above for Compound C, thereby producing Compound D where R is  $-CO_2Et$ , which after conversion to a compound of Z may be reacted with a compound having the Formula II to thereby produce a corresponding compound having the  
30 Formula I. The  $-CO_2Et$  group may be removed by standard procedures.

The syntheses of the starting compounds represented by Formula II are illustrated in the following schemes.


-25-

Scheme 1 below illustrates the formation of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-alkyl-4-oxo-3-quinolinecarboxylic acid.



-26-

In Scheme 1 above the 2-pentafluorophenyl-4,4-dimethyl-2-oxazoline III is reacted with alkyl lithium at -20°C to +25°C to give the 2-(2,3,4,5-tetrafluoro-6-alkylphenyl)-4,4-dimethyl-2-oxazoline IV which is hydrolyzed under acidic conditions (preferably refluxing dilute hydrochloric acid) to give the corresponding benzoic acid V. Compound V is reacted with oxalyl chloride and the product condensed with the dianion of monoethyl malonate (prepared from monoethyl malonic acid and n-butyl lithium in THF) to produce ketoester VII. This ketoester is treated with triethyl orthoformate in acetic anhydride to form adduct VIII. Reaction of compound VIII with cyclopropylamine in t-butanol or ether gives enamine IX; other primary amines can be used in this reaction, such as aliphatic amines (ethylamine etc.) and aromatic amines (p-fluoroaniline, 2,4-difluoroaniline, etc.) The enamine is reacted with potassium t-butoxide in dry t-butanol to form the desired cyclized compound X, which can be hydrolyzed in refluxing acid to give Compound XI.



-28-

In Scheme 2 above the acid XII is converted to its acid chloride via reaction with oxalyl chloride, and the acid chloride is treated with 2-amino-2-methyl-1-propanol to give N-(2-hydroxy-1,1-dimethylethyl)-2,4,5-trifluorocarboxamide (Compound XIII). This amide is cyclized to the crucial intermediate oxazoline XIV by reaction with thionyl chloride in chloroform. Compound XIV is then treated with a base, preferably lithium diisopropylamide, in THF or ether at -78°C and quenched with trimethylsilyl chloride to produce silylated oxazoline XV. Compound XV is treated with base (again, preferably lithium diisopropylamide) in THF or ether at 0°-20°C and then quenched with an alkyl iodide to give, upon work-up, the alkylated intermediate XVI. Removal of the trimethylsilyl group is accomplished by treatment with cesium fluoride in wet DMF; the resulting compound XVII is hydrolyzed to the corresponding benzoic acid XVIII in refluxing dilute hydrochloric acid. This benzoic acid is elaborated into 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-alkyl-4-oxo-3-quinolinecarboxylic acid using the methodology previously described in Scheme I.


Alternatively, silylated intermediate XVI can be transformed into a variety of 3-substituted compounds via ipso attack on the trimethylsilyl group. For example, Compound XVI is reacted with chlorine in the presence of iron powder and then hydrolyzed in dilute refluxing acid to give 3-chloro-2,4,5-trifluoro-6-alkylbenzoic acid XIX; this acid is elaborated as before to give quinoline XX. Similarly, oxazoline XVI is treated with N-bromosuccinimide in chloroform (or with pyridinium bromide perbromide in dichloromethane) to give the analogous 3-bromo oxazoline which is hydrolyzed and carried on to give Compound XXII. Reaction of intermediate XVI with lead tetraacetate

-29-

and trifluoroacetic acid, followed by acid hydrolysis, gives 2,4,5-trifluoro-3-hydroxy-6-alkylbenzoic acid (XXIII); the phenol can be converted to the methyl ether via reaction with methyl iodide and potassium carbonate in acetone. This 2,4,5-trifluoro-3-methoxy-6-alkylbenzoic acid is carried on to the 8-methoxy quinoline XXIV (where R = CH<sub>3</sub>); further treatment with HBr cleaves the methyl ether to give the corresponding 8-hydroxy quinoline XXIV (where R = H). Finally, nitration of silylated compound XVI with nitric acid in sulfuric acid and hydrolysis of the consequent compound yields nitro acid XXV, which is further elaborated to afford 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-alkyl-8-nitro-4-oxo-3-quinolinecarboxylic acid XXVI (where R<sub>1</sub> = O). Reduction of the nitro group to the amino group can be accomplished using Raney nickel to yield quinoline XXVI (where R<sub>1</sub> = H).

-30-

Scheme 2A below outlines an alternative route to the 5-alkyl,8-chloro quinolones.



In Scheme 2A above the oxazoline XIV is treated with a base, preferably lithium diisopropylamide, in THF at  $-78^\circ\text{C}$  and quenched with hexachloro acetone to produce the chloro oxazoline XXXVIII. Compound XXXVIII is treated again with a base, preferably lithium diisopropylamide, in THF at  $0^\circ\text{C}$  and




-31-

quenched with an alkyl iodide to give, upon work-up, the intermediate XXXIX. Hydrolysis of the oxazoline moiety in refluxing dilute hydrochloric acid gives benzoic acid XIX. This acid is elaborated into

5 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-alkyl-4-oxo-3-quinolinecarboxylic acid (XX).

-32-

Scheme 3 below illustrates synthesis of 5,8-dialkyl quinolines.




- 5 In Scheme 3 above oxazoline XIV (prepared in Scheme II) is treated with a base, preferably lithium diisopropylamide, in THF at -78°C and is quenched with an alkyl halide (such as methyl iodide, ethyl iodide, etc.) to give Compound XXVII, where R = alkyl.
- 10 Treatment with additional base (preferably lithium diisopropylamide) in ether at 0°C followed by addition of an alkyl halide affords dialkyl oxazolines such as XXVIII. The intermediates are hydrolyzed and carried

-33-

on as before to give 5,8-dialkyl-1-cyclopropyl-6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acids XXX.

-34-

Scheme 4 below illustrates a synthesis of 5-alkyl naphthyridines



-35-


In Scheme 4 above pyridine ester XXXI (Chem. Pharm. Bull. 35 (1987), p 2280) is reacted with a base such as lithium diisopropylamide in THF at low temperature followed by an alkyl halide such as ethyl iodide or methyl iodide; hydrolysis of the ester in  
5 dilute acid affords compound XXXIII.

Alternatively, ester XXXI can be hydrolyzed in dilute acid to give pyridine acid XXXV which is, in turn, converted to the corresponding oxazoline in the  
10 usual manner (see Scheme 2). This oxazoline (Compound XXXVI) is reacted with an alkyl lithium (such as methyl lithium), then rearomatized with DDQ or chloranil. This sequence of reactions gives the alkyl-substituted pyridine XXXVII, which yields, upon  
15 acid hydrolysis, the necessary intermediate XXXIII.

Compound XXXIII can be elaborated to the 5-alkyl-7-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid in the usual manner.


-36-

Scheme 5



-37-

Also prepared by this method:




8

In Scheme 5 above, the known naphthyridine acid 1 (US Patent 4,663,457, 1987) is reacted with oxalyl chloride and DMF and then quenched with absolute ethanol to give ester 2. Reduction of the double bond is accomplished with sodium cyanoborohydride to afford compound 3, which is then treated with sec-butyllithium at -78°C. This dianion is treated with methyl iodide to give the alkylated intermediate 4. The double bond is reintroduced in a series of steps: first, treatment with sodium hydride, followed by addition of phenylselenenyl chloride and oxidation with hydrogen peroxide. The final ester 5 can then be reacted with a variety of amines in the usual fashion.

-38-

Scheme 6 outlines the synthesis of 5-alkyl, 8-trifluoromethyl derivatives.





-39-

Scheme 6 begins with the treatment of 2,4,5-tri-fluorobromobenzene with a base, preferably lithium diisopropylamide, in THF at -78°C. This anion is quenched with carbon dioxide to give, upon  
5 acidification, acid XL. The acid is reacted with HF/SF<sub>4</sub> at 120°C to afford the trifluoromethyl derivative XLI. The requisite acid functionality is introduced via halogen-metal exchange (preferably with butyl lithium in ether at -78°C) followed by carbon  
10 dioxide quench and acidification. Compound XLII is then treated with oxalyl chloride to form the acid chloride and added to 2-amino-2-methyl-1-propanol in chloroform at 0°C to produce hydroxy amide XLIII. Cyclization to the key intermediate oxazoline XLIX is  
15 accomplished in the usual manner - that is, treatment with thionyl chloride followed by sodium hydride. Deprotonation of XLIV by lithium diisopropylamide at -78°C and reaction of the anion with an alkyl iodide yields the fully substituted oxazoline XLV which can  
20 be elaborated into the target quinolone XLVI using the previously established methodology.

The compounds of the invention are capable of forming both pharmaceutically acceptable acid addition and/or base salts. Base salts are formed with metals  
25 or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzyl-ethylenediamine, chlorprocaine, choline,  
30 diethanolamine, ethylenediamine, N-methylglucamine, and procaine.

Pharmaceutically acceptable acid addition salts are formed with organic and inorganic acids.

Examples of suitable acids for salt formation are  
35 hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, gluconic, fumaric,

-40-

succinic, ascorbic, maleic, methanesulfonic, and the like. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce either a mono or di, etc salt in the conventional manner. The free base forms may be regenerated by treating the salt form with a base. For example, dilute solutions of aqueous base may be utilized. Dilute aqueous sodium hydroxide, potassium carbonate, ammonia, and sodium bicarbonate solutions are suitable for this purpose. The free base forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but the salts are otherwise equivalent to their respective free base forms for purposes of the invention. Use of excess base where R' is hydrogen gives the corresponding basic salt.

The compound of the invention can exist in unsolvated as well as solvated forms, including hydrated forms. In general, the solvated forms, including hydrated forms and the like are equivalent to the unsolvated forms for purposes of the invention.

The alkyl groups contemplated by the invention comprise both straight and branched carbon chains of from one to about six carbon atoms. Representative of such groups are methyl, ethyl, propyl, isopropyl, and the like.

The cycloalkyl groups contemplated by the invention comprise those having three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

The alkoxy groups contemplated by the invention comprise both straight and branched carbon chains of from one to about six carbon atoms unless otherwise specified. Representative of such groups are methoxy, ethoxy, propoxy, *i*-propoxy, *t*-butoxy, hexoxy, and the like.

-41-

The term, haloalkyl, is intended to include halogen substituted straight and branched carbon chains of from two to four carbon atoms. Those skilled in the art will recognize that the halogen substituent may not be present on the  $\alpha$ -carbon atom of the chain. Representative of such groups are  $\beta$ -fluoroethyl,  $\beta$ -chloroethyl,  $\beta,\beta$ -dichloroethyl,  $\beta$ -chloropropyl,  $\beta$ -chloro-2-propyl,  $\gamma$ -iodobutyl, and the like.

10 The term halogen is intended to include fluorine, chlorine, bromine, and iodine unless otherwise specified.

Certain compounds of the invention may exist in optically active forms. The pure D isomer, pure L isomer as well as mixtures thereof; including the racemic mixtures, are contemplated by the invention. Additional asymmetric carbon atoms may be present in a substituent such as an alkyl group. All such isomers as well as mixtures thereof are intended to be included in the invention.

20 The compounds of the invention can be prepared and administered in a wide variety of oral and parenteral dosage forms. It will be obvious to those skilled in the art that the following dosage forms may comprise as the active component, either a compound of Formula I or a corresponding pharmaceutically acceptable salt of a compound of Formula I.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories. A solid carrier can be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, or tablet disintegrating agents; it

-42-

can also be an encapsulating material. In powders, the carrier is a finely divided solid which is in admixture with the finely divided active compound. In the tablet the active compound is mixed with carrier  
5 having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from 5 or 10 to about 70 percent of the active ingredient. Suitable solid carriers are magnesium  
10 carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation  
15 of the active compound with encapsulating material as carrier providing a capsule in which the active component (with or without other carriers) is surrounded by carrier, which is thus in association with it. Similarly, cachets are included. Tablets,  
20 powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

Liquid form preparations include solutions, suspensions, and emulsions. As an example may be mentioned water or water-propylene glycol solutions  
25 for parenteral injection. Such solutions are prepared so as to be acceptable to biological systems (isotonicity, pH, etc). Liquid preparations can also be formulated in solution in aqueous polyethylene glycol solution. Aqueous solutions suitable for oral  
30 use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents as desired. Aqueous suspension suitable for oral use can be made by dispersing the finely divided active component in  
35 water with viscous material, i.e., natural or synthetic gums, resins, methyl cellulose, sodium

-43-

carboxymethyl cellulose, and other well known suspending agents.

Preferably, the pharmaceutical preparation is in unit dosage form. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself or it can be the appropriate number of any of these packaged forms.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from 1 mg to 100 mg according to the particular application and the potency of the active ingredient.

In therapeutic use as agents for treating bacterial infections the compounds utilized in the pharmaceutical method of this invention are administered at the initial dosage of about 3 mg to about 40 mg per kilogram daily. A daily dose range of about 6 mg to about 14 mg per kilogram is preferred. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The compounds of the invention display anti-bacterial activity when tested by the microtitration

-44-

dilution method as described in Heifetz, et al, Antimicrob. Agents & Chemoth., 6, 124 (1974), which is incorporated herein by reference. By use of this method, the following minimum inhibitory concentration values (MICs in  $\mu\text{g/ml}$ ) were obtained for

5 representative compounds of the invention.

-45-

## IN VITRO ANTIBACTERIAL ACTIVITY

Minimal Inhibitory Concentration  
MIC ( $\mu\text{g/ml}$ )

Organisms	Compound Ex. 1	Compound Ex. 2	Compound Ex. 3	Compound Ex. 4	Compound Ex. 5
Enterobacter cloacae MA 2646	0.025	0.006	0.1	0.05	0.025
Escherichia coli Vogel	0.025	0.006	0.05	0.05	0.05
Klebsiella pneumoniae MGH-2	0.025	0.013	0.1	0.1	0.05
Providencia rettgeri M 1771	0.05	0.025	0.2	0.2	0.10
Pseudomonas aeruginosa UI-18	0.2	0.4	0.8	0.8	0.4
Staphylococcus aureus H 228	0.025	0.013	0.013	0.006	0.025
Staphylococcus aureus UC-76	0.025	0.003	0.003	0.003	0.013
Streptococcus faecalis MGH-2	0.05	0.025	0.025	0.025	0.10
Streptococcus pneumoniae SV-1	0.025	0.003	0.003	0.003	0.025
Streptococcus pyogenes C-203	0.05	0.013	0.006	0.003	0.006

## IN VITRO ANTIBACTERIAL ACTIVITY

Minimal Inhibitory Concentration  
MIC ( $\mu\text{g/ml}$ )

Organisms	Compound Ex. 6	Compound Ex. 7	Compound Ex. 8	Compound Ex. 9	Compound Ex. 10	Compound Ex. 11
Enterobacter cloacae MA 2646	0.013	0.025	0.05	0.10	0.05	0.05
Escherichia coli Vogel	0.013	0.013	0.025	0.10	0.025	0.013
Klebsiella pneumoniae MGH-2	0.05	0.05	0.10	0.40	0.10	0.05
Proteus rettgeri M 1771	0.10	0.20	0.10	0.80	0.20	0.05
Pseudomonas aeruginosa UI-18	0.20	0.80	0.40	1.6	0.20	0.20
Staphylococcus aureus H 228	0.10	0.10	0.10	0.025	0.05	0.025
Staphylococcus aureus UC-76	0.025	0.025	0.025	0.006	0.025	0.013
Streptococcus faecalis MGH-2	0.05	0.10	0.40	0.10	0.20	0.05
Streptococcus pneumoniae SV-1	0.05	0.025	0.013	0.006	0.10	0.025
Streptococcus pyogenes C-203	0.10	0.05	0.025	0.013	0.20	0.05



-47-

IN VITRO ANTIBACTERIAL ACTIVITY  
Minimal Inhibitory Concentration  
MIC ( $\mu\text{g/ml}$ )

Organisms	Compound			Compound			Compound		
	Ex. 12	Ex. 13	Ex. 14	Ex. 14a	Ex. 15	Ex. 15	Ex. 14a	Ex. 15	Ex. 15
Enterobacter cloacae MA 2646	0.05	0.05	0.1	0.4	0.025	0.025	0.4	0.025	0.025
Escherichia coli Vogel	0.05	0.05	0.05	0.4	0.1	0.1	0.4	0.025	0.025
Klebsiella pneumoniae MGH-2	0.20	0.2	0.4	1.6	0.2	0.2	1.6	0.1	0.1
Proteus rettgeri M 1771	0.40	0.4	0.4	3.1	0.2	0.2	3.1	0.2	0.2
Pseudomonas aeruginosa UI-18	1.6	1.6	0.8	3.1	0.8	0.8	3.1	0.8	0.8
Staphylococcus aureus H 228	0.8	0.4	0.2	0.4	0.05	0.05	0.4	0.05	0.05
Staphylococcus aureus UC-76	0.2	0.1	0.05	0.2	0.025	0.025	0.2	0.025	0.025
Streptococcus faecalis MGH-2	3.1	0.8	0.4	1.6	0.2	0.2	1.6	0.2	0.2
Streptococcus pneumoniae SV-1	6.3	1.6	0.4	0.8	0.1	0.1	0.8	0.1	0.1
Streptococcus pyogenes C-203	12.5	3.1	0.4	1.6	0.1	0.1	1.6	0.1	0.1

-48-

## IN VITRO ANTIBACTERIAL ACTIVITY

Minimal Inhibitory Concentration  
MIC (µg/ml)

Organisms	Compound Ex. 17	Compound Ex. 17a	Compound Ex. 17b	Compound Ex. 17c	Compound Ex. 17e	Compound Ex. 18	Compound Ex. 19
Enterobacter cloacae MA 2646	0.05	0.2	0.4	0.05	0.4	0.013	0.025
Escherichia coli Vogel	0.05	0.1	0.2	0.05	0.4	0.013	0.013
Klebsiella pneumoniae MGH-2	0.1	0.4	0.8	0.1	0.8	0.025	0.05
Proteus rettgeri M 1771	0.2	0.8	1.6	0.2	0.8	0.1	0.2
Pseudomonas aeruginosa UI-18	0.8	1.6	1.6	0.8	3.1	0.2	0.4
Staphylococcus aureus H 228	0.1	0.1	0.1	0.05	0.05	0.25	0.2
Staphylococcus aureus UC-76	0.025	0.025	0.05	0.013	0.013	0.013	0.1
Streptococcus faecalis MGH-2	0.2	0.4	0.8	0.1	0.2	0.05	0.2
Streptococcus pneumoniae SV-1	0.05	0.05	0.05	0.025	0.013	0.013	0.1
Streptococcus pyogenes C-203	0.1	0.1	0.1	0.05	0.025	0.025	0.4

-49-

The following nonlimiting examples illustrate the inventors' preferred methods for preparing the compounds of the invention.

#### PREPARATION OF STARTING MATERIALS

##### 5 Example A

##### 2-(2,3,4,5-Tetrafluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline

A solution of 21.2 g (80.0 mmol) of 2-(pentafluoro-phenyl)-4,4-dimethyl-2-oxazoline (Bull. Chem. Soc. Jpn., 57, 225 (1984)) in 300 ml of dry ether was cooled to -20°C under argon and treated with 60 ml of 1.6M methyl lithium (96.0 mmol). The solution was stirred at -20°C for two hours, then stirred at room temperature overnight. The mixture was diluted with water, and the organic layer was dried over magnesium sulfate and concentrated to give 20.8 g of the title compound as an orange oil.

##### Example B

##### 2,3,4,5-Tetrafluoro-6-methylbenzoic acid

20 A mixture of 20.5 g (73.4 mmol) of 2-(2,3,4,5-tetrafluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline in 200 ml of 6N hydrochloric acid was refluxed for 18 hours, then cooled to room temperature. The solution was extracted with ethyl acetate. The  
25 extract was washed with water, dried over magnesium sulfate, and concentrated. The residue was suspended in water which was made basic (pH 11) with 1M sodium hydroxide and was extracted with ether; the aqueous phase was acidified (pH 2) with 1N hydrochloric acid  
30 and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and concentrated to give 8.4 g of the title compound as a tan solid, mp 80-82°C.

-50-

## Example C

2,3,4,5-Tetrafluoro-6-methylbenzoyl chloride

5 A solution of 8.2 g (39.4 mmol) of 2,3,4,5-tetrafluoro-6-methylbenzoic acid, 6.0 g (47.2 mmol) of oxalyl chloride, and 100 ml of dichloromethane was treated with three drops of DMF. The solution was stirred for three hours, then concentrated to give 8.8 g of the title compound as a yellow liquid. The product was used as is in the next step.

10

## Example D

Ethyl 3-(2,3,4,5-tetrafluoro-6-methylphenyl)- $\beta$ -oxo-propanoate

15 A solution of 10.1 g (76.5 mmol) of malonic acid monoethylester, bipyridyl (catalytic), and 200 ml of dry THF was cooled to -35°C under argon, treated with 52 ml of 1.5M n-butyllithium (78 mmol), and warmed to -5°C. To this mixture was added 52 ml of 1.5M n-butyllithium (78 mmol) until a pale pink color persisted for 10 minutes. The suspension was cooled  
20 to -78°C and was treated with a solution of 8.8 g (38.8 mmol) of 2,3,4,5-tetrafluoro-6-methylbenzoyl chloride in 100 ml of dry THF. The reaction mixture was stirred at -78°C for 45 minutes, then warmed to -35°C and poured into a mixture of ice and 1N  
25 hydrochloric acid (77 ml). The organic layer was washed with 5% sodium bicarbonate solution, 3M hydrochloric acid, and water and dried over magnesium sulfate. Concentration gave an orange oil which was chromatographed on silica gel (E. Merck 230-400 Mesh),  
30 eluting with 80:20 chloroform:ethyl acetate, to give 8.2 g of the title compound.

-51-

## Example E

Ethyl 2-(2,3,4,5-Tetrafluoro-6-methyl-benzoyl)-3-ethoxyacrylate

5 A solution of 8.1 g (29.1 mmol) of ethyl 3-(2,3,4,5-tetrafluoro-6-methylphenyl)- $\beta$ -oxo-propanoate, 7.2 g (43.3 mmol) of triethyl orthoformate, and 70 ml of acetic anhydride was refluxed for 3.5 hours. The solution was cooled to room temperature and concentrated under high vacuum to  
10 give 9.1 g of the title compound. The product was used as is in the next step.

## Example F

Ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-cyclopropylaminoacrylate

15 To a solution of 9.0 g (27.0 mmol) of ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-ethoxyacrylate in 30 ml of absolute ethanol at 5°C was added 1.68 g (29.4 mmol) of cyclopropylamine. The mixture was  
20 stirred at 5°C for 1.5 hours and at room temperature for 2.5 hours. The solution was concentrated to an oil which was triturated with hexane to give a tan solid. The crude product was recrystallized from hexane to give 9.07 g of the title compound, mp 72-74°C.

## Example G

Ethyl 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate

30 To a mixture of 9.05 g (26.3 mmol) of ethyl 2-(2,3,4,5-tetrafluoro-5-methylbenzoyl)-3-cyclopropylaminoacrylate in 100 ml of dry t-butanol was added a slurry of 3.25 g (29.0 mmol) of potassium t-butoxide in 20 ml of dry t-butanol, and the mixture was stirred at 60°C for four hours. The suspension was cooled to room temperature and concentrated to a paste which was

-52-

partitioned between dichloromethane and 1N hydrochloric acid. The organic layer was separated, dried over magnesium sulfate, and concentrated. Recrystallization from ethyl acetate:hexane gave  
5 4.70 g of the title compound, mp 176-177°C.

## Example H

1-Cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 4.6 g (14.1 mmol) of ethyl 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate in 100 ml of 6M hydrochloric acid was refluxed for four hours. The solution was cooled to room temperature and the solids were filtered, washed with water, and dried to give 3.9 g of the  
15 title compound, mp 234-235°C.

In a similar manner, 1-cyclopropyl-5-ethyl-6,7,8-trifluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid and 1,5-dicyclopropyl-6,7,8-trifluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid were prepared.

## 20 Example I

N-(2-Hydroxy-1,1-dimethylethyl)-2,4,5-trifluorobenzamide

A solution of 19.4 g (110 mmol) of 2,4,5-trifluorobenzoic acid (JP 58,150,543 (Cl. C07C69) Sept. 7, 1983). 15.2 g (120 mmol) of oxalyl chloride and 250 ml of dichloromethane was treated with four drops of DMF, and the mixture was stirred at room temperature for four hours. The mixture was concentrated to a oil and was redissolved in 100 ml of  
30 dichloromethane. This solution was added dropwise to a solution of 19.6 g (240 mmol) of 3-amino-2-methyl-1-propanol in 200 ml of dichloromethane at 5°C, and the reaction mixture was stirred at room temperature overnight. The solids were filtered, and the filtrate

-53-

was washed with 5% sodium bicarbonate, 1N hydrochloric acid, and water. The organic layer was dried over magnesium sulfate and concentrated to give 24.5 g of the title compound, mp 114-116°C.

5

## Example J

2-(2,4,5-Trifluorophenyl)-4,4-dimethyl-2-oxazoline

To a solution of 24.4 g (98.7 mmol) of N-(2-hydroxy-1,1-dimethylethyl)-2,4,5-trifluorocarbox-  
amide in 200 ml of chloroform was added 25 ml  
10 (342 mmol) of thionyl chloride dropwise. The solution was stirred overnight at room temperature, then concentrated by half. The mixture was diluted with ether, and the solid was removed by filtration. This solid was dissolved in water, made basic (pH 8) with  
15 10% sodium hydroxide, and extracted with ethyl acetate. The organic layer was dried over magnesium sulfate and concentrated to give 19.0 g of the title compound, mp 53-54°C.

## Example K

20 2-(2,4,5-Trifluoro-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline

A solution of 8.7 ml (62.1 mmol) of diisopropyl-amine in 100 ml of dry THF under argon was cooled to -78°C and treated with 28.3 ml (56.6 mmol) of 2.0M  
25 n-butyllithium. The LDA solution was stirred at -78°C for 15 minutes. To this solution was added a solution of 11.8 g (51.5 mmol) of 2-(2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline in 50 ml of THF, and the reaction mixture was stirred for one hour at -78°C.  
30 To the reaction mixture was added 13 ml (102.5 mmol) of chlorotrimethylsilane, and the solution was warmed to room temperature. Water was added; the organic layer was dried over magnesium sulfate and

-54-

concentrated. The crude product was chromatographed on silica gel (E. Merck 230-400 Mesh), eluting with 80:20 chloroform:ethyl acetate to give 12.9 g of the title compound, mp 71-72°C.

5

## Example L

2-(2,4,5-Trifluoro-6-methyl-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline

A solution of 0.64 ml (4.57 mmol) of diisopropylamine in 20 ml of dry THF under argon was cooled to -78°C and treated with 2.1 ml (4.20 mmol) of 2.0N n-butyllithium. The LDA solution was stirred at -78°C for 15 minutes, then warmed to 0°C. To this solution was added a solution of 1.05 g (3.5 mmol) of 2-(2,4,5-trifluoro-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline in 5 ml of THF; the reaction mixture was stirred at 0°C for 45 minutes, then quenched with 1.50 g (10.6 mmol) of methyl iodide. The solution was stirred at room temperature for three hours and diluted with water. The organic layer was washed with water, dried over magnesium sulfate, and concentrated to give 1.00 g of the title compound as an oil.

In a similar manner, 1-cyclopropyl-5-ethyl-6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid and 1-cyclopropyl-6,7-difluoro-1,4-dihydro-4-oxo-5-ethyl-3-quinolinecarboxylic acid were prepared.

Alternatively, the trimethylsilyl group was displaced with chlorine (Chem. Abstr. 54, 20932 (1960)) or with bromine (J. Am. Chem. Soc. 70, 433 (1948)), and the oxazoline was hydrolyzed to give 3-chloro-2,4,5-trifluoro-6-methylbenzoic acid and 3-bromo-2,4,5-trifluoro-6-methylbenzoic acid, respectively. These intermediates were elaborated into 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid 8-bromo-1-



-55-

cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid.

In addition, the trimethylsilyl group was reacted with lead tetraacetate/trifluoroacetic acid to  
5 introduce a hydroxyl group (Tet. Lett. 10, 853 (1974)) and was also reacted with nitric acid to introduce a nitro group (J. Chem. Soc. 498 (1957)).

Following the usual procedures, the following compounds were prepared: 1-cyclopropyl-6,7-difluoro-  
10 1,4-dihydro-8-hydroxy-5-methyl-4-oxo-3-quinolinecarboxylic acid; 1-cyclopropyl-6,7-difluoro-1,4-dihydro-8-methoxy-5-methyl-4-oxo-3-quinolinecarboxylic acid; 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-8-nitro-4-oxo-3-quinolinecarboxylic acid, and  
15 8-amino-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid.

#### Example M

##### 7-Chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid

20 Ethyl 2,6-dichloro-5-fluoronicotinate (Chem. Pharm. Bull. 35(6), 2280 (1987)) was treated with lithium diisopropylamide and quenched with methyl iodide to give, upon work-up, ethyl 2,6-dichloro-5-fluoro-4-methylnicotinate. This material was  
25 hydrolyzed to give the corresponding acid which was elaborated in the usual manner to give 7-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid. 7-chloro-1-cyclopropyl-5-ethyl-6-fluoro-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid was synthesized in the  
30 same manner.

-56-

## Example N

2,4,5-Trifluoro-3,6-dimethylbenzoic acid

The 2-(2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline was also treated with lithium diisopropyl-  
5 amide followed by methyl iodide to give  
2-(2,4,5-trifluoro-3-methylphenyl)-4,4-dimethyl-2-oxazoline. This intermediate was, in turn, treated  
with lithium diisopropylamide, then with methyl  
10 iodide, to give 2-(2,4,5-trifluoro-3,6-dimethylphenyl)-  
4,4-dimethyl-2-oxazoline. Hydrolysis of the oxazoline  
gave 2,4,5-trifluoro-3,6-dimethylbenzoic acid, which  
was elaborated into 1-cyclopropyl-6,7-difluoro-1,4-  
dihydro-5,8-dimethyl-4-oxo-3-quinolinecarboxylic acid  
in the usual manner.

15

## Example O

2-(2,4,5-Trifluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline

A solution of 12.0 g (38.0 mmol) of 2-[2,4,5-trifluoro-6-methyl-3-(trimethylsilyl)phenyl]-4,4-  
20 dimethyl-2-oxazoline, 5.85 g (38.5 mmol) of cesium  
fluoride, 110 ml of dimethylformamide, and 15 ml of  
water was stirred for 18 hours at room temperature.  
The reaction mixture was poured into water and  
25 extracted with ethyl acetate; the organic phase was  
washed with water, dried over magnesium sulfate, and  
concentrated to give 9.1 g of liquid.

## Example P

2-(3-Chloro-2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline

30 A solution of 7.6 ml (54.2 mmol) of diisopropyl-  
amine in 100 ml of dry THF was cooled to -78°C under  
argon, treated with 20.5 ml (47.2 mmol) of 2.3M  
n-butyllithium, and stirred for 15 minutes. To this  
solution was added a solution of 10.3 g (45.0 mmol) of

-57-

2-(2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline in 100 ml of dry THF. The reaction mixture was stirred at -78°C for 45 minutes. To this mixture was added 26.5 g (100 mmol) of hexachloroacetone, and the solution was warmed to room temperature. Water was added; the organic phase was washed with water, 1N hydrochloric acid, and 5% sodium bicarbonate, and was dried over magnesium sulfate. Concentration gave a dark oil which was chromatographed on silica gel to give 7.05 g of the title compound as a yellow oil.

## Example Q

2-(3-Chloro-2,4,5-trifluoro-6-methyl-4,4-dimethyl-2-oxazoline

A solution of 5.5 ml (39.2 mmol) of diisopropylamine in 125 ml of dry THF was cooled to -78°C under argon, treated with 13.8 ml (31.7 mmol) of 2.3M n-butyllithium, and stirred for 15 minutes. To this solution was added a solution of 7.00 g (26.5 mmol) of 2-(3-chloro-2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline in 75 ml of dry THF. The mixture was stirred at -78°C for 30 minutes and at 0°C for 60 minutes. To this solution was added 11.3 g (79.6 mmol) of methyl iodide, and the mixture was stirred at room temperature overnight. Water was added; the organic phase washed with 1N HCl, 5% sodium bicarbonate, and water. The solution was dried over magnesium sulfate and concentrated to an oil which was chromatographed on silica gel to give 6.3 g of clear orange oil.

## Example R

2,4,5-Trifluoro-6-methylbenzoic acid

A mixture of 9.1 g (37.4 mmol) 2-(2,4,5-trifluoro-6-methylphenyl)-4,4-dimethyl-2-oxazoline in 200 ml of 6M hydrochloric acid was refluxed overnight, then

-58-

cooled to room temperature. The solution was extracted with ethyl acetate, and the extract was washed with water, dried over magnesium sulfate, and concentrated. The residue was suspended in water  
5 which was made basic (pH 11) with 1N NaOH, washed with ether, and acidified (pH 2) with 1N HCl. The solution was extracted with ethyl acetate. The organic phase was washed with water, dried over magnesium sulfate, and concentrated to give 5.8 g of the title compound,  
10 mp 108-110°C.

## Example S

3-Chloro-2,4,5-trifluoro-6-methylbenzoic acid

As in Example D, the title compound was prepared from 2-(3-chloro-2,4,5-trifluoro-6-methylphenyl)-4,4-  
15 dimethyl-2-oxazoline and 6N hydrochloric acid. The desired acid was obtained as a tan solid, mp 104-106°C.

## Example T

2,4,5-Trifluoro-6-methylbenzoyl chloride

20 A solution of 5.8 g (30.5 mmol) of 2,4,5-trifluoro-6-methylbenzoic acid, 4.7 g (37.0 mmol) of oxalyl chloride, and 100 ml of dichloromethane was treated with three drops of DMF. The reaction mixture was stirred at room temperature for two hours, then  
25 concentrated to give 6.3 g of the title compound as an oily solid. The product was used "as is" in the next step.

## Example U

3-Chloro-2,4,5-Trifluoro-6-methylbenzoyl chloride

30 The title compound was prepared from 3-chloro-2,4,5-trifluoro-6-methylbenzoic acid and oxalyl chloride following the same procedure used in Example F.

-59-

## Example V

Ethyl 3-(2,4,5-trifluoro-6-methylbenzoyl)- $\beta$ -oxo-propanoate

A solution of 8.0 g (60.5 mmol) of malonic acid monoethylester, bipyridyl (catalytic) and 200 ml of dry THF was cooled to -35°C under argon, treated with 32 ml of 1.9M n-butyllithium (60.8 mmol), and warmed to -5°C. To this suspension was added another 32 ml of 1.9M n-butyllithium until a pale pink color persisted for 10 minutes. The mixture was cooled to -78°C. To this mixture was added a solution of 6.3 g (30.2 mmol) of 2,4,5-trifluoro-6-methylbenzoyl chloride in 75 ml of dry THF, and the reaction mixture was stirred at -78°C for one hour. The solution was then warmed to -35°C, poured onto a mixture of ice and 1N hydrochloric acid (70 ml), and extracted with ethyl acetate. The organic layer was washed with 5% sodium bicarbonate, 3M hydrochloric acid, and water, and was stirred over magnesium sulfate. Concentration gave an orange oil which was chromatographed on silica gel (E. Merck 230-400 Mesh), eluting with 80:20 chloroform:ethyl acetate, to give 7.2 g of the title compound.

## Example W

Ethyl 3-(3-chloro-2,4,5-trifluoro-6-methylbenzoyl)- $\beta$ -oxopropanoate

The procedure outlined for Example H was used to prepare the title compound from the dianion of malonic acid monoethyl ester and 3-chloro-2,4,5-trifluoro-6-methylbenzoyl chloride. The crude product was also chromatographed on silica gel to give the desired product as an orange oil.

-60-

## Example X

Ethyl 2-(2,4,5-trifluoro-6-methylbenzoyl)-3-ethoxy-acrylate

A solution of 7.1 g (27 mmol) of ethyl  
5 (3-(2,4,5-trifluoro-6-methylbenzoyl)- $\beta$ -oxo-propanoate,  
6.8 g (41 mmol) of triethyl orthoformate and 60 ml of  
acetic anhydride was refluxed for three hours, cooled  
to room temperature, and concentrated to give 8.4 g of  
the title compound. The crude material was used as is  
10 in the next step.

## Example Y

Ethyl 2-(3-chloro-2,4,5-trifluoro-6-methylbenzoyl)-3-ethoxy acrylate

The procedure outlined in Example J was followed  
15 to prepare the title compound from ethyl 3-(3-chloro-  
2,4,5-trifluoro-6-methylbenzoyl)- $\beta$ -oxo-propanoate,  
triethyl orthoformate, and acetic anhydride.

## Example Z

Ethyl 2-(2,4,5-Trifluoro-6-methylbenzoyl)-3-cyclo-propylaminoacrylate

To a solution of 8.3 g (26 mmol) of ethyl  
2-(2,4,5-trifluoro-6-methylbenzoyl)-3-ethoxyacrylate  
in 30 ml of absolute ethanol at 5°C was added 1.64 g  
(29 mmol) of cyclopropylamine. The reaction mixture  
25 was stirred at 5°C for 90 minutes and at room  
temperature for two hours. The solution was  
concentrated to give a brown oil which was dissolved  
in hexane and reconcentrated to give a tan solid.  
Recrystallization from hexane gave 7.2 g of colorless  
30 crystals, mp 69-72°C.

The following compounds were prepared in  
identical fashion from the appropriate ethoxyacrylate:

-61-

- a) Ethyl 2-(3-chloro-2,4,5-trifluoro-6-methylbenzoyl)-3-cyclopropylaminoacrylate, mp 77-80°C;
- b) Ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-ethylamino acrylate, hygroscopic solid;
- 5 c) Ethyl 3-(2,4-difluoroanilino)-2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)acrylate, viscous oil;
- d) Ethyl 3-(2-bromoethylamino)-2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)acrylate, mp 95-100°C;
- e) Ethyl 3-(ethylamino)-2-(2,4,5-trifluoro-6-methylbenzoyl)acrylate, hygroscopic solid;
- 10 f) Ethyl 3-(2,4-difluoroanilino)-2-(2,4,5-trifluoro-6-methylbenzoyl)acrylate, mp 79-83°C; and
- g) Ethyl 3-(2-bromoethylamino)-2-(2,4,5-trifluoro-6-methylbenzoyl)acrylate.

15 Example AA

Ethyl 1-cyclopropyl-6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylate

A solution of 7.2 g (22 mmol) of ethyl 2-(2,4,5-trifluoro-5-methylbenzoyl)-3-cyclopropylaminoacrylate  
20 in 100 ml of dry t-butanol was treated portionwise with 2.8 g (25 mmol) of potassium t-butoxide, and the reaction mixture was stirred at 60°C for five hours. The suspension was cooled to room temperature and concentrated. The residue was partitioned between  
25 dichloromethane and 1N hydrochloric acid; the organic phase was washed with water, dried over magnesium sulfate, and concentrated. The crude product was slurried in boiling ethanol, filtered, and air-dried to give 4.2 g of the title compound.

30 The following compounds were prepared in a similar fashion and purified as noted:

a) Ethyl 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-

-62-

carboxylate, mp 151-153°C (chromatographed on silica gel);

b) Ethyl 1-ethyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate, mp 185-187°C (recrystallized from ethyl acetate);

c) Ethyl 1-(2-bromoethyl)-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate, mp 149-150°C (recrystallized from ethyl acetate hexane).

d) Ethyl 1-ethyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate, mp 189-191°C.

#### Example BB

Ethyl 6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-1-vinyl-3-quinolinecarboxylate

A rapidly stirred suspension of 1.98 g (5.08 mmol) of ethyl 1-(2-bromoethyl)-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate, 3.50 g (25.3 mmol) of ground potassium carbonate, and 40 ml of DMF was heated at 80°C under argon for four hours. The suspension was concentrated and the residue was partitioned between methylene chloride and water. The organic layer was dried over magnesium sulfate and concentrated to give 1.52 g of the title compound as a DMF complex, mp 150-152°C.

#### Example CC

Ethyl 6,7,8-trifluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate

To a cold (5°C) solution of 2.77 g (6.64 mmol) of ethyl 3-(2,4-difluoroanilino)-2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)acrylate in 60 ml of dry THF was added 0.32 g of 60% sodium hydride. The solution was stirred overnight at room temperature, then concentrated to an orange foam. The residue was partitioned between methylene chloride and 1N HCl.



-63-

The organic phase was washed with water, dried over magnesium sulfate, and concentrated to an orange solid which was recrystallized (ethyl acetate:hexane) to give 1.55 g of the title compound, mp 152-154°C.

5

## Example DD

Ethyl 6,7-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate

The procedure outlined in Example AA was used to prepare the title compound from ethyl 3-(2,4-difluoro-anilino)-2-(2,4,5-trifluoro-6-methylbenzoyl)acrylate, mp 161-164°C.

10

## Example EE

1-Cyclopropyl-6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylic acid

A suspension of 4.1 g (13.3 mmol) of ethyl 1-cyclopropyl-6,7-difluoro-1,4-dihydro-4-oxo-3-quinolinecarboxylate in 150 ml of 6N hydrochloric acid was refluxed for six hours, then cooled to room temperature. The solids were filtered, washed with water and ether, and dried to give 3.2 g of the title compound, mp >300°C.

15

20

The following compounds were prepared in a similar fashion:

a) 1-Ethyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 199-201°C;

25

b) 1-Ethyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp >300°C;

c) 8-Chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,

mp 212-214°C.

30

-64-

## Example FF

7-Chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester

7-Chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid, (U.S. Patent 4,663,457) (20.0 g, 71 mmol) and dimethylformamide (0.5 ml) were added to dichloromethane (750 ml) to give a tan slurry. Oxalyl chloride (7.4 ml, 85 mmol) was added to this slurry over one minute and the reaction mixture stirred for 90 minutes, then an additional 2.0 ml of oxalyl chloride was added and stirring continued for 60 minutes. To the resulting brown solution was added absolute ethanol (4.3 ml, 78 mmol) and the mixture stirred for four hours and then cooled to 0°C and stored overnight. The reaction was warmed to room temperature and an additional 2 ml of absolute ethanol was added and the stirring continued for three hours. The reaction was evaporated to a brown solid. The solid was heated in THF, filtered, and cooled to 0°C. The crystals formed were collected and dried to give the title compound, (11.1 g, 50%).

## Example GG

7-Chloro-1-cyclopropyl-6-fluoro-1,2,3,4-tetrahydro-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester, 3

In absolute ethanol (200 ml) was suspended the compound prepared in Example FF (3.0 g, 9.6 mmol) and sodium cyanoborohydride (0.7 g, 10 mmol) and three drops of concentrated HCl was added, giving a bright yellow solution. As the reaction progressed and was monitored by TLC (silica gel, CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 9:1 v/v) additional aliquots of concentrated HCl were added as needed to maintain the progress of the reaction. After six hours the reaction was quenched by adding it to 300 ml of water. The mixture was

-65-

extracted several times with  $\text{CH}_2\text{Cl}_2$  and the combined organic layers dried, filtered, and evaporated to a yellow solid. This solid was filtered through silica gel with  $\text{CH}_2\text{Cl}_2$  and after evaporation the solid was  
5 crystallized from isopropyl ether. The collected crystals were further purified by column chromatography on silica gel with  $\text{CH}_2\text{Cl}_2$  to give the title compound (2.2 g, 73%).

10 The following compound was prepared in the same manner:

a) 7-chloro-6-fluoro-1-(2,4-difluorophenyl)-1,2,3,4-tetrahydro-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester

Example HH

15 7-Chloro-1-cyclopropyl-6-fluoro-2,3,4-tetrahydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester

Compound GG (4.5 g, 14 mmol) was dissolved in THF (170 ml) and cooled to  $-70^\circ\text{C}$ . Then sec-butyl lithium (22.2 ml, 28 mmol, 1.3M) was added dropwise over  
20 30 minutes, always keeping the internal temperature  $-70^\circ\text{C}$ . After stirring at  $-70^\circ\text{C}$  for one hour, methyl iodide (0.9 ml, 14 mmol) was added and the reaction stirred at  $-70^\circ\text{C}$  for seven hours. The reaction flask  
25 was transferred to a Dewar containing dry ice/isopropanol and allowed to stand for 17 hours. At the end of this time period the reaction temperature had warmed to  $-25^\circ\text{C}$ . The reaction was quenched by the addition of saturated  $\text{NH}_4\text{Cl}$  solution (50 ml) and  
30 diluted with an equal volume of  $\text{CH}_2\text{Cl}_2$ . The organic layer was separated and washed with saturated NaCl solution, dried, filtered, and evaporated to an oil. This oil was purified by column chromatography on silica gel with  $\text{CH}_2\text{Cl}_2$  to give, after combining and

-66-

evaporating the appropriate fraction, the title compound (3.91 g, 85%).

## Example II

5 7-Chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester

Using the procedure of Reich, et al, J. Amer. Chem. Soc., (1975) 97, 5434, the compound prepared in Example HH (0.68 g, 2.1 mmol) was converted into the title compound (0.44 g, 64%). Purification was  
10 achieved by crystallization from isopropyl ether.

## Example JJ

3-Bromo-2,5,6-trifluorobenzoic acid

n-Butyl lithium (2.6 M in hexanes, 32 ml, 84 mmol) was added over 10 minutes to a solution of  
15 diisopropylamine (8.89 g, 88 mmol) in THF (80 ml) stirred under N<sub>2</sub> at 0°C. After a further 10 minutes at 0°, the solution was transferred by catheter over 40 minutes to a solution of 2,4,5-trifluorobromobenzene (16.88 g, 80 mmol) in THF (200 ml) stirred  
20 under N<sub>2</sub> at -78°C. After a further 15 minutes the solution was blown through a catheter over ~2 minutes onto a slurry of CO<sub>2</sub> (~200 ml) in ether (400 ml) with vigorous stirring. When the CO<sub>2</sub> evaporated the slurry was washed with dilute HCl (1 M, 200 ml) and water  
25 (100 ml). The organic phase was extracted with dilute NaOH (0.5 M, 2x100 ml). The aqueous phase was extracted with ether (2x100 ml), and the combined organic phases were washed with water (100 ml), saturated brine (100 ml), and dried (MgSO<sub>4</sub>). The  
30 solvent was removed under reduced pressure to give 3-bromo-2,5,6-trifluorobenzoic acid (17.25 g, 84.5%) as white microcrystalline needles; mp 114-6°C (sublimation).

-67-

## Example KK

1-Bromo-2,4,5-trifluoro-3-(trifluoromethyl)benzene

3-Bromo-2,5,6-trifluorobenzoic acid (16.92 g, 66 mmol) was heated with SF<sub>4</sub> (60 g) and HF (30 g) in a stainless steel bomb at 120°C for 8 hours. When the reaction cooled to 25°C, the volatiles were vented through KOH traps, and when gas evolution ceased the vessel was extracted with CH<sub>2</sub>Cl<sub>2</sub> (150 ml). This solution was washed with diluted NaHCO<sub>3</sub> solution (saturated/2, 50 ml), saturated brine (50 ml), and dried (MgSO<sub>4</sub>). The solvent was removed by distillation through a 15-cm Vigreux column, and the residue was distilled under N<sub>2</sub> through a shortpath stillhead at 147-150°C to give 1-bromo-2,4,5-trifluoro-3-(trifluoromethyl)benzene (15.79 g, 83%) as a pale yellow oil. nmr (CDCl<sub>3</sub>) δ 7.67 (1H, d of t, J<sub>d</sub> = 6 Hz, J<sub>t</sub> 8.1 Hz, aromatic).

## Example LL

2,4,5-Trifluoro-3-(trifluoromethyl)benzoic acid

A solution of n-butyl lithium (2.6 M in hexanes, 9.6 ml, 25 mmol) was added dropwise through an addition funnel over 15 minutes to a solution of 1-bromo-2,4,5-trifluoro-3-(trifluoromethyl)benzene (7.00 g, 25 mmol) in ether (100 ml) stirred under N<sub>2</sub> at -78°C. After 5 minutes the mixture was rapidly blown by catheter onto a suspension of dry ice (100 g) in ether (100 ml). After 5 minutes TFA (2 ml) was added to this. When the solution had warmed up to 20°C, it was washed with diluted HCl (0.5 M, 20 ml), and extracted with dilute base (0.5 N, 2x50 ml). The combined basic extracts were washed with ether (25 ml), made acidic with concentrated HCl (~4 ml), and extracted with ether (3x50 ml). The combined ethereal extracts were washed with water (50 ml), saturated brine (50 ml), and dried (MgSO<sub>4</sub>). The

-68-

solvent was removed under reduced pressure to give 2,4,5-trifluoro-3-(trifluoromethyl)benzoic acid (4.21 g, 69%) as white microscopic needles; mp 87-90°C. Nmr (CDCl<sub>3</sub>) δ 11.80 (1H, br s, OH), 8.05 (1H, d of t, J<sub>d</sub> = 6 Hz, J<sub>t</sub> = 9 Hz, aromatic).

## Example MM

N-(2-Hydroxy-1,1-dimethylethyl)-2,4,5-trifluoro-3-trifluoromethyl)benzamide

A solution of 4.88 g (20.0 mmol) of 2,4,5-trifluoro-3-(trifluoromethyl)benzoic acid, 2.80 g (22.0 mmol) of oxalyl chloride, and 50 ml of methylene chloride was treated with 1 drop of DMF and stirred at room temperature for 4 hours. The solution was concentrated to a yellow oil which was dissolved in methylene chloride (20 ml) and added to a cold (ice bath) solution of 2.53 g (25 mmol) of triethylamine, 1.96 g (22 mmol) of 2-amino-2-methyl-1-propanol, and 40 ml of methylene chloride. The mixture was allowed to warm slowly to room temperature overnight. The solution was poured into 50 ml of 1 N HCl, and the organic layer was separated and washed with water. The solution was dried over magnesium sulfate and concentrated to give 5.81 g of the title compound as a yellow oil.

25

## Example NN

2-[2,4,5-Trifluoro-3-(trifluoromethyl)phenyl]-4,4-dimethyl-2-oxazoline

A solution of 5.81 g (18.4 mmol) of N-(2-hydroxy-1,1-dimethylethyl)-2,4,5-trifluoro-3-(trifluoromethyl)benzamide in 100 ml of chloroform at 0°C was treated dropwise with 5 ml of thionyl chloride. The mixture was allowed to warm to room temperature overnight. The solution was concentrated to a yellow oil which was dissolved in 20 ml of DMF and treated with 0.8 g

-69-

(21.6 mmol) of 60% sodium hydride. This reaction mixture was stirred at room temperature for 18 hours, then poured into 50 ml of dilute  $\text{NaHCO}_3$ . The solution was extracted with ethyl acetate; the organic phase was washed with water and dried over magnesium sulfate. Concentration in vacuo gave an orange oil which was chromatographed on silica, eluting with 2% methanol in chloroform, to give 2.62 g of a yellow oil.

10

## Example OO

2-2,4,5-Trifluoro-3-(trifluoromethyl)-6-methylphenyl]-4,4-dimethyl-2-oxazoline

A solution of 1.12 g (11.0 mmol) of diisopropylamine in 5 ml of THF was cooled to 0°C under nitrogen, treated with 4.0 ml of 2.5 M n-butyllithium, and stirred for 10 minutes. This lithium diisopropylamide solution was added dropwise to a solution of 2.36 g (8 mmol) of 2-[2,4,5-trifluoro-3-(trifluoromethyl)-phenyl]-4,4-dimethyl-2-oxazoline in 5 ml of THF at -78°C. The solution was stirred at -78°C for one hour, then quenched with 2.24 g (16 mmol) of methyl iodide. The mixture was allowed to warm slowly to room temperature, stirred for one hour, and poured into 10 ml of 1 N HCl. This solution was extracted with ether, and the extract was washed with water, dried over magnesium sulfate, and concentrated to give 2.28 g of the title compound.

30

## Example PP

3-(Exo-amino)-8-azabicyclo[3.2.1]octane, dihydro-chloride

A mixture of 4.6 g (20 mmole) of 8-(phenylmethyl)-8-azabicyclo[3.2.1]octan-3-one, oxime [J. R. Bagley and T. N. Riley, J. Heterocyclic Chem., 19, 485 (1982)], 0.5 g of 10% rhodium on carbon, and 100 ml of

-70-

acetic acid was hydrogenated until the requisite amount of hydrogen was taken up. The reaction mixture was filtered and two equivalents of HCl was added. The solid was filtered to yield 2.80 g of the title compound, mp >300°C.

## Example QQ

3(Endo-amino)-8-azabicyclo[3.2.1]octane, dihydro-chloride

A solution of 7.33 g (25 mmol) of 3-(endo-amino)-8-(phenylmethyl)-8-azabicyclo[3.2.1]octane dihydrochloride [P. Dostert et al, Eur. J. Med. Chem.-Chim. Ther., 19, 105 (1984)], 1.0 g of 20% palladium on carbon and 100 ml of methanol was hydrogenated until the required amount of hydrogen was taken up. The reaction mixture was filtered and the filtrate was evaporated to 4.5 g of the title compound which was used without purification.

## Example 1

1-Cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A suspension of 0.85 g (2.85 mmol) of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 1.00 g (11.6 mmol) of anhydrous piperazine, and 20 ml of acetonitrile was refluxed for five hours, then stirred at room temperature overnight. The precipitate was filtered, washed with water and acetonitrile, and dried to give 0.91 g of the title compound, mp 205-206°C.

## Example 2

7-(3-Amino-1-pyrrolidinyl)-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 1.00 g (3.36 mmol) of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-



-71-

carboxylic acid, 0.63 g (3.38 mmol) of 3-(t-butoxy-carbonyl)aminopyrrolidine, 1.00 g (9.91 mmol) of triethylamine, and 35 ml of acetonitrile was refluxed for five hours, then stirred at room temperature overnight. The precipitate was filtered and washed with acetonitrile and ether. The crude product was suspended in 20 ml of 6M hydrochloric acid and 20 ml of glacial acetic acid and was heated at 60°C for two hours. The solution was concentrated to an oil which was triturated with isopropanol. The solid was filtered and washed with ether to give 1.04 g of the title compound as the hydrochloride salt, mp >300°C.

## Example 3

1-Cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-carboxylic acid

A mixture of 0.80 g (2.70 mmol) of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-carboxylic acid, 0.41 g (3.20 mmol) of N-ethyl-3-pyrrolidinemethanamine, 0.82 g (8.10 mmol) of triethylamine, and 25 ml of acetonitrile was refluxed for four hours, then stirred at room temperature overnight. The precipitate was filtered, washed with acetonitrile and ether, and dried to give 0.90 g of the title compound, mp 198-199°C.

## Example 4

7-[3-(Aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.60 g (2.02 mmol) of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-carboxylic acid, 0.28 g (2.45 mmol) of 3-methyl-3-pyrrolidinemethanamine, 0.61 g (6.06 mmol) of triethylamine, and 20 ml of acetonitrile was refluxed

-72-

for four hours, then stirred at room temperature overnight. The precipitate was filtered, washed with ether, and dried to give 0.61 g of the title compound, mp 182-184°C.

5

## Example 5

1-Cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-7-[3-methyl-1-piperazinyl]-4-oxo-3-quinolinecarboxylic acid

A suspension of 0.80 g (2.69 mmol) of 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 1.08 g (10.8 mmol) of 2-methylpiperazine, and 20 ml of acetonitrile was refluxed for three hours, then cooled in an ice bath. The precipitate was filtered, washed with water and acetonitrile, and dried to give 0.76 g of the title compound, mp 187-188°C.

## Example 6

1-Cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A suspension of 0.70 g (2.50 mmol) of 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.86 g (10.0 mmol) of anhydrous piperazine and 20 ml of acetonitrile was refluxed for five hours, then stirred at room temperature overnight. The precipitate was filtered, washed with water and acetonitrile, and dried to give 0.85 g of the title compound, mp 226-228°C.

## Example 7

1-Cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.75 g (2.68 mmol) of 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 1.07 g (10.4 mmol) of 2-methylpiperazine and 30 ml of acetonitrile was refluxed for

-73-

five hours, then stirred at room temperature overnight. The precipitate was filtered, washed with water/ethanol and acetonitrile, and dried to give 0.42 g of the title compound, mp 189-192°C.

5

## Example 8

7-(3-Amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.70 g (2.50 mmol) of 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-  
10 carboxylic acid, 0.56 g (3.00 mmol) of 3-(t-butoxy-carbonyl)aminopyrrolidine, 0.76 g (7.52 mmol) of triethylamine, and 25 ml of acetonitrile was refluxed for 4.5 hours, then stirred at room temperature overnight. The solids were filtered and washed with  
15 acetonitrile and ether. The crude product was dissolved in 20 ml of 6N hydrochloric acid and 20 ml of acetic acid and was stirred at room temperature for three hours. The solution was concentrated to an oil which was triturated with 2:1 ether:isopropanol. The  
20 solids were filtered and washed with ether to give 0.95 g of the title compound as the hydrochloride salt, mp >300°C.

## Example 9

25 7-[3-(Aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-carboxylic acid

A mixture of 0.61 g (2.18 mmol) of 1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinoline-  
30 carboxylic acid, 0.56 g (2.61 mmol) of 3-[(t-butoxy-carbonyl)aminomethyl]-3-methylpyrrolidine, 0.66 g (6.54 mmol) of triethylamine, and 25 ml of acetonitrile was refluxed for six hours, then stirred overnight at room temperature. The precipitate was filtered and washed with acetonitrile and ether. The

-74-

crude product was suspended in 20 ml of 6N hydrochloric acid and 20 ml of glacial acetic acid and was stirred at room temperature for three hours. The solution was concentrated and the residue was  
5 trituated with ether. The solid was filtered and washed with ether to give 0.61 g of the title compound as the hydrochloride, mp 250-252°C.

## Example 10

10 8-Chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A suspension of 0.38 g (1.21 mmol) of 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.42 g (4.88 mmol) of piperazine and 20 ml of acetonitrile was refluxed for  
15 four hours, then stirred at room temperature overnight. The precipitate was filtered and washed with water and acetonitrile to give 0.32 g of the title compound, mp 234-235°C.

## Example 11

20 7-(3-Amino-1-pyrrolidinyl)-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.50 g (1.60 mmol) of 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.36 g (1.93 mmol) of  
25 3-(t-butoxycarbonyl)aminopyrrolidine, 0.48 g (4.75 mmol) of triethylamine, and 20 ml of acetonitrile was refluxed for four hours, then stirred at room temperature overnight. The solution was  
30 concentrated and the residue was trituated with ether:hexane (1:1) and filtered. The solid was washed with water and hexane. The crude product was suspended in 15 ml of dichloromethane and 1.5 ml of trifluoroacetic acid and was stirred at room

-75-

temperature for four hours. The solution was concentrated to a gold solid which was suspended in water, made basic (pH 11) with 10% sodium hydroxide, and filtered. The solution was then neutralized (pH 7.10), and the precipitate was filtered and washed with water to give 0.29 g of the title compound, mp 124-126°C.

## Example 12

10 1-Ethyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A mixture of 0.45 g (1.58 mmol) of 1-ethyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.54 g (6.27 mmol) of anhydrous piperazine, and 20 ml of acetonitrile was refluxed for three hours, then cooled to room temperature. The solids were filtered and washed with water, acetonitrile, and ether to give 0.48 g of the title compound, mp 223-225°C.

## Example 13

20 7-(3-Amino-1-pyrrolidinyl)-1-ethyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.36 g (1.25 mmol) of 1-ethyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.26 g (1.39 mmol) of 3-(t-butoxycarbonyl)aminopyrrolidine, 0.38 g (3.76 mmol) of triethylamine, and 20 ml of acetonitrile was refluxed for five hours, then stirred at room temperature overnight. The solids were filtered and washed with acetonitrile and ether. The crude product was dissolved in 5 ml of 6N hydrochloric acid and 5 ml of glacial acetic acid and stirred for five hours at room temperature. The solution was concentrated to a solid which was suspended in water, made basic (pH 12), filtered through a fiberglass pad, and neutralized

-76-

(pH 6.8). The solids were filtered and washed with water to give 0.32 g of white solid, mp 218-220°C.

## Example 14

5 6,8-Difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A solution of 0.43 g (1.08 mmol) of ethyl 6,7,8-trifluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.37 g  
10 (4.30 mmol) of anhydrous piperazine, and 20 ml of acetonitrile was refluxed overnight, cooled to room temperature, and concentrated. The residue was taken up in 10 ml of 6N hydrochloric acid and refluxed for two hours. The mixture was cooled and the solids were  
15 filtered. The crude product was suspended in water which was made basic (pH 12), filtered through a fiberglass pad, and neutralized (pH 6.5). The solids were filtered and washed with water and ether to give 0.37 g of the title compound, mp 283-284°C.

20 The following compound was prepared following the same procedure:

a) 6,8-Difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid, mp 240-242°C.

25 Example 15

7-(3-Amino-1-pyrrolidinyl)-6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A solution of 0.40 g (1.00 mmol) of ethyl 6,7,8-trifluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.22 g (1.18 mmol) of  
30 3-t-butoxycarbonylaminopyrrolidine, 0.30 g (3.00 mmol) of triethylamine, and 15 ml of acetonitrile was

-77-

refluxed for 18 hours. The mixture was cooled and concentrated. The residue was dissolved in 10 ml of 6N hydrochloric acid, refluxed for three hours, and cooled to room temperature. The solids were filtered, washed with water and ether, and suspended in water. The suspension was made basic (pH 12) and filtered through a fiberglass pad, and the filtrate was neutralized to pH 6.7. The solids were filtered and washed with water and ether to give 0.38 g of the title compound, mp 230-232°C.

## Example 16

6,8-Difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid

A solution of 0.69 g (1.80 mmol) of ethyl 6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-1-vinyl-3-quinolinecarboxylate, 0.62 g (7.2 mmol) of anhydrous piperazine, and 20 mmol of acetonitrile was refluxed for 18 hours, cooled, and concentrated. The residue was suspended in 25 ml of 1N sodium hydroxide and heated at 80°C for 90 minutes. The clear yellow solution was cooled to room temperature, filtered, and neutralized (pH 6.8) with 6N hydrochloric acid. The solids were filtered, washed with water and ether, and dried to give 0.32 g of the title compound, mp 222-225°C.

The following compound was prepared in identical fashion:

a) 6,8-Difluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid, mp 232-235°C.

-78-

## Example 17

6-Fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid

A solution of 0.76 g (2.00 mmol) of ethyl  
5 6,7-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylate, 0.69 g  
(8.00 mmol) of anhydrous piperazine, and 30 ml of acetonitrile was refluxed for 18 hours, cooled, and concentrated. The residue was dissolved in 20 ml 6N  
10 hydrochloric acid and refluxed for three hours. The suspension was cooled, concentrated by half and filtered, and the solids were washed with water. The crude product was suspended in water which was made basic (pH 12), filtered, and neutralized to pH 6.8.  
15 The precipitate was filtered and neutralized to pH 6.8. The precipitate was filtered, washed with water, and dried to give 0.58 g of the title compound, mp 198-200°C.

The following compounds were also prepared by  
20 following essentially the same procedure:

- a) 6-Fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid, mp 188-191°C;
- b) 6-Fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-  
25 5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid, mp 213-215°C;
- c) 7-(3-Amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 232-234°C;
- 30 d) 7-[3-(Ethylamino)methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 196-198°C; and
- e) 7-[3-(Aminomethyl)-3-methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-  
35 oxo-3-quinolinecarboxylic acid, mp 181-184°C.



-79-

## Example 18

7-(3-Amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid

5        Triethylamine (0.17 ml, 1.2 mmol), 3-(1,1-dimethyl-ethoxycarbonylamino)-pyrrolidine (0.22 g, 2.1 mmol) and 0.39 g (1.2 mmol) of 7-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester were dissolved in  
10        acetonitrile (10 ml) and the mixture heated to reflux for four hours, then cooled and diluted with ether (50 ml). This solution was washed with saturated solutions of  $\text{KHCO}_3$  and  $\text{NaCl}$  and dried over  $\text{Na}_2\text{SO}_4$ . It was necessary to add  $\text{CH}_2\text{Cl}_2$  to maintain a homogeneous  
15        solution. After filtration and evaporation the product was dissolved in ether and allowed to stand. The crystals formed were collected to give 0.56 g of the intermediate ester. The intermediate ester was dissolved in acetic acid (15 ml) and 6N  $\text{HCl}$  (1 ml) was  
20        added and the mixture heated to reflux for two hours, then evaporated to a gum. This gum was dissolved in ethanol (10 ml), and 5N  $\text{NaOH}$  (2 ml) was added and the mixture stirred for two hours. The reaction was evaporated to a gum and dissolved in water (60 ml) to  
25        give a solution at pH 12. The pH was adjusted to 6.5 and the solid formed collected and washed with water and dried to give the title compound (0.38 g).

## Example 19

1-Cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-1,8-naphthyridine-3-carboxylic acid

30        The procedure used in Example 18 was employed to prepare the title compound in 58% yield.

-80-

## Example 20

1-Ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-1-(piperazinyl)-3-quinolinecarboxylic acid

A suspension of 0.67 g (2.50 mmol) of 1-ethyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, 0.86 g (10.0 mmol) of anhydrous piperazine, and 25 ml of acetonitrile was refluxed for six hours, then cooled to room temperature. The solids were filtered, washed with water and ether, and dried to give 0.58 g of the title compound, mp 225-227°C.

The following compounds were prepared by following essentially the same procedure:

a) 1-Ethyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 180-182°C.

b) 7-(3-Amino-1-pyrrolidinyl)-1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 210-213°C.

c) 1-Ethyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid, mp 228-231°C.

d) 1-Ethyl-6-fluoro-1,4-dihydro-5-methyl-7-3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid, mp 219-221°C.

e) 7-[3-(Aminomethyl)-3-methyl-1-pyrrolidinyl]-1-ethyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid, mp 223-225°C.

## Example 21

7-[3-(Endo-amino)-8-azabicyclo[3.2.1]oct-8-yl]-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid

A mixture of 0.50 g (1.6 mmol) of 8-chloro-1-cyclopropyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-3-

-81-

quinolinecarboxylic acid, 0.36 g (1.8 mmol) of  
3-(endo-amino)-8-azabicyclo[3.2.1]octane  
dihydrochloride, 0.72 ml (4.8 mmol) of 1,8-diazabicyclo-  
[5.4.0]undec-7-ene and 15 ml of acetonitrile was  
5 heated at reflux for 18 hours. The suspension was  
cooled to room temperature, diluted with ether, and  
refrigerated. The resulting solid was filtered,  
washed with ethanol and ether, and dried to give the  
title compound.

10 The following compounds were prepared in  
identical fashion:


Example a. 7-[3-(endo-amino)-8-azabicyclo[3.2.1]-  
oct-8-yl]-6,8-difluoro-1-(2,4-difluorophenyl)-1,4-  
dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid.

15 Example b. 7-[3-(endo-amino)-8-azabicyclo[3.2.1]-  
oct-8-yl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-  
methyl-4-oxo-3-quinolinecarboxylic acid.







-82-

## CLAIMS

1. A compound of formula



or a pharmaceutically acceptable acid addition or base salt thereof wherein Z is



-83-

5        wherein  $R_4$  is hydrogen, alkyl of from one to  
four carbon atoms, a cycloalkyl of from three to  
six carbon atoms,  $R'$  is hydrogen, hydroxyl, alkyl  
of from one to four carbon atoms, phenyl or  
phenyl substituted by halogen, alkyl or alkoxy,  $n$   
10        is an integer of from 0 to 4,  $R_5$  and  $R_6$  are each  
independently hydrogen, lower alkyl or  
cycloalkyl;

$X$  is  $CH$ ,  $CF$ ,  $CCl$ ,  $CBr$ ,  $N$ ,  $CCF_3$ ,  $CNH_2$ ,  $CNO_2$ ,  $CR$ ,  
or  $COR'$  wherein  $R$  is lower alkyl and  $R'$  is  
hydrogen or lower alkyl;

15         $R_3$  is lower straight, branched, or cyclic alkyl  
of from one to three carbon atoms;


20         $R_2$  is alkyl of from one to four carbon atoms,  
vinyl, haloalkyl, hydroxyalkyl of from two to  
four carbon atoms, cycloalkyl of from three to  
six carbon atoms, phenyl or phenyl substituted by  
halogen, alkyl,  $NH_2$  or  $OH$ ; and

$R_1$  is hydrogen, alkyl of from one to six carbon  
atoms, or a cation.

2. A compound according to Claim 1, wherein  $X$  is  $CH$ ,  
 $CF$ ,  $CCl$  or  $N$ .
3. A compound according to Claim 1, wherein  $R_2$  is  
cyclopropyl, ethyl or 2,4-difluorophenyl.
4. A compound according to Claim 1, wherein  $R_3$   
methyl, ethyl, isopropyl, or cyclopropyl.

-84-











5. A compound according to Claim 1, wherein  $R_1$  is hydrogen or a pharmaceutically acceptable base salt thereof.
6. A compound according to Claim 1, wherein Z is



wherein  $R_4$  is hydrogen, methyl or cyclopropyl,  $R'$  is hydrogen or methyl,  $n$  is 0, 1, or 2,  $R_5$  and  $R_6$  are each independently hydrogen or methyl.

7. A compound according to Claim 6 wherein Z is selected from the group consisting of:

-85-



or

-86-

8. A compound according to Claim 1 and selected from the groups consisting of:

- 5 1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
10 1-cyclopropyl-7-[3-[(ethylamino)methyl]-1-pyrrolidinyl]-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
15 1-cyclopropyl-6,8-difluoro-1,4-dihydro-5-methyl-7-[3-methyl-1-piperazinyl]-4-oxo-3-quinolinecarboxylic acid,  
1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
20 1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
25 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
30 8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,  
35




-87-

- 1-ethyl-6,7-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-1-ethyl-6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 5 6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,
- 10 6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-6,8-difluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 15 6,8-difluoro-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid,  
6,8-difluoro-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-1-vinyl-3-quinolinecarboxylic acid,
- 20 6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid,  
6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3-methyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,
- 25 6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-7-(3,5-dimethyl-1-piperazinyl)-4-oxo-3-quinolinecarboxylic acid,  
7-(3-amino-1-pyrrolidinyl)-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 30 7-[3-(ethylamino)methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid,
- 35 7-[3-(aminomethyl)-3-methyl-1-pyrrolidinyl]-6-fluoro-1-(2,4-difluorophenyl)-1,4-dihydro-5-

-88-

methyl-4-oxo-3-quinolinecarboxylic acid, and pharmaceutically acceptable salts thereof.

9. A compound according to Claim 1 and selected from the group consisting of 1-cyclopropyl-7-(3-amino-1-pyrrolidinyl)-6-fluoro-5-methyl-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester and 1-cyclopropyl-7-(piperazinyl)-6-fluoro-5-methyl-1,4-dihydro-4-oxo-1,8-naphthyridine-3-carboxylic acid ethyl ester.
10. A compound according to Claim 1 named 7-[3-(endo-amino)-8-azabicyclo[3.2.1]oct-8-yl]-8-chloro-1-cyclopropyl-6-fluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid.
11. A pharmaceutical composition comprising an antibacterially effective amount of compound according to Claim 1 together with a pharmaceutically acceptable carrier.
12. A method of treating bacterial infections in mammals which comprises administering to said mammal a pharmaceutical composition according to Claim 11.
13. A process for the preparation of a compound according to Claim 1 which comprises reacting a compound of formula




-89-

5 wherein L is fluorine or chlorine with an amine  
corresponding to Z as defined in Claim 1 and, if  
desired, converting the resulting product to a  
pharmaceutically acceptable acid addition or base  
salt thereof by known means.

14. A compound named 1-cyclopropyl-6,7,8-trifluoro-1,4-dihydro-5-methyl-4-oxo-3-quinolinecarboxylic acid ethyl ester.
15. A compound named 2,3,4,5-tetrafluoro-6-methylbenzoic acid.
16. A compound named 2,3,4,5-tetrafluoro-6-methylbenzoyl chloride.
17. A compound named ethyl 3-(2,3,4,5-tetrafluoro-6-methylphenyl)- $\beta$ -oxo-propanoate.
18. A compound named ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-ethoxyacrylate.
19. A compound named ethyl 2-(2,3,4,5-tetrafluoro-6-methylbenzoyl)-3-cyclopropylaminoacrylate.
20. A compound named 2-(2,4,5-trifluoro-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline.
21. A compound named 2-(2,4,5-trifluoro-6-methyl-3-trimethylsilylphenyl)-4,4-dimethyl-2-oxazoline.
22. A compound named 2-(2,4,5-trifluorophenyl)-4,4-dimethyl-2-oxazoline.


-90-

23. A process for the preparation of compound of formula




wherein R is alkyl which comprises reacting a pentafluorooxazoline with alkyl lithium producing a compound of formula

5




followed by acidic hydrolysis.

24. A process for the preparation of compounds of formula




wherein R is alkyl which comprises  
(a) reacting a compound of formula


-91-



- 5 with a base and trimethylsilyl chloride producing a compound of formula



- (b) reacting that compound with a base and an alkylhalide producing a compound of formula




- 10 (c) removing the  $\text{SiMe}_3$ , and  
(d) hydrolyzing the resulting compound.

-92-


25. A process for the preparation of compounds of Formula I by

(a) reacting a compound of formula




5

with oxalyl chloride and dimethylformamide and quenching with alcohol to produce the corresponding ester




(b) reducing the double bond to produce a compound of formula



10

(c) treating the compound from step (b) with a base, then methyl iodide to produce the alkylated compound



(d) reintroducing the double bond and reacting the resulting naphthyridine with the desired amine by known means.

26. Use of a compound according to Claims 1 to 10 for the preparation of antibacterial pharmaceuticals.
27. Process for the preparation of antibacterial pharmaceuticals comprising the incorporation of a compound according to Claims 1 to 10 into a pharmaceutically acceptable carrier.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>4</sup> : C07D 215/56, 401/04, 471/04 C07D 451/04, A61K 31/44 A61K 31/47, C07C 51/00 // (C07D 471/04, 221:00, 221:00)</p>	<p>A3</p>	<p>(11) International Publication Number: <b>WO 89/ 06649</b></p> <p>(43) International Publication Date: 27 July 1989 (27.07.89)</p>
<p>(21) International Application Number: PCT/US89/00278</p> <p>(22) International Filing Date: 23 January 1989 (23.01.89)</p> <p>(31) Priority Application Numbers: 147,462 280,924</p> <p>(32) Priority Dates: 25 January 1988 (25.01.88) 9 December 1988 (09.12.88)</p> <p>(33) Priority Country: US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 280,924 (CIP) Filed on 9 December 1988 (09.12.88)</p> <p>(71) Applicant (for all designated States except US): WARN- ER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US).</p>	<p>(72) Inventors; and (75) Inventors/Applicants (for US only) : DOMAGALA, John, Michael [US/US]; 776 Georgetown, Canton, MI 48188 (US). HAGEN, Susan, Elizabeth [US/US]; 1035 Paddington Road, Canton, MI 48187 (US). KIELY, John, Steven [US/US]; 4138 Sunset Court, Ann Arbor, MI 48103 (US).</p> <p>(74) Agents: ANDERSON, Elizabeth, M.; Warner-Lambert Company, 2800 Plymouth Road, Ann Arbor, MI 48105 (US) et al.</p> <p>(81) Designated States: AT, AT (European patent), AU, BE (European patent), CH (European patent), DE, DE (European patent), DK, FI, FR (European patent), GB, GB (European patent), IT (European patent), JP, KR, LU, LU (European patent), NL, NL (European patent), NO, SE, SE (European patent), US.</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 28 December 1989 (28.12.89)</p>	
<p>(54) Title: ANTIBACTERIAL AGENTS</p> <p>(57) Abstract</p> <p>Novel naphthyridine-, and quinolinecarboxylic acids as antibacterial agents are described as well as methods for their manufacture, formulation, and use in treating bacterial infections including the description of certain novel intermediates used in the manufacture of the antibacterial agents.</p>		



***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria  
AU Australia  
BB Barbados  
BE Belgium  
BG Bulgaria  
BJ Benin  
BR Brazil  
CF Central African Republic  
CG Congo  
CH Switzerland  
CM Cameroon  
DE Germany, Federal Republic of  
DK Denmark  
FI Finland

FR France  
GA Gabon  
GB United Kingdom  
HU Hungary  
IT Italy  
JP Japan  
KP Democratic People's Republic  
of Korea  
KR Republic of Korea  
LI Liechtenstein  
LK Sri Lanka  
LU Luxembourg  
MC Monaco  
MG Madagascar

ML Mali  
MR Mauritania  
MW Malawi  
NL Netherlands  
NO Norway  
RO Romania  
SD Sudan  
SE Sweden  
SN Senegal  
SU Soviet Union  
TD Chad  
TG Togo  
US United States of America

# INTERNATIONAL SEARCH REPORT

International Application No PCT/US 89/00278

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC <sup>4</sup> C 07 D 215/56, C 07 D 401/04, C 07 D 471/04, C 07 D 451/04, IPC: A 61 K 31/44, A 61 K 31/47, C 07 C 51/00, // ./.		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System <sup>1</sup>	Classification Symbols	
IPC <sup>4</sup>	C 07 D 215/00, C 07 D 401/00, C 07 D 471/00, C 07 C 45/00, C 07 C 51/00, C 07 C 63/00	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched *		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT *</b>		
Category *	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	US, A, 4341784 (J. MATSUMOTO et al.) 27 July 1982, see the abstract cited in the application --	1-3,5-7,11, 13,26,27
A	EP, A, 0207420 (DAIICHI SEIYAKU CO. LTD) 7 January 1987, see claims 1,7,11,21 --	1-3,5,6,11, 13,26,27
A	EP, A, 0242789 (DAINIPPON PHARMACEUTICAL CO., LTD) 28 October 1987, see claims 1,2,12,14,15 --	1-3,5,6,11, 13,26,27
A	US, A, 4146719 (T. IRIKURA) 27 March 1979, see claim 1; abstract cited in the application --	1-3,5-7,11, 26,27
A	US, A, 4649144 (J. MATSUMOTO et al.) 10 March 1987, see claims 1,2,12 cited in the application --	1-3,5,6,11, 26,27
. / .		
<p>* Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
5th October 1989	27 NOV 1989	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	T.K. WILLIS	

# INTERNATIONAL SEARCH REPORT PCT/US 89/00278

International Application No -2-

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC <sup>4</sup> : (C 07 D 471/04, 221:00, 221:00)		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
IPC <sup>4</sup>		
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	US, A, 4571396 (M.P. HUTT et al.) 18 February 1986, see claims 1,3-5,31 cited in the application --	1-3,5,11, 26,27
A	Journal of Medicinal Chemistry, vol. 23, 1980, American Chemical Society, H. Koga et al.: "Structure-activity relationships of antibacterial 6,7- and 7,8-disubstituted 1-alkyl-1,4- dihydro-4-oxoquinoline-3-carboxylic acids", pages 1358-1363, see page 1358, abstract; page 1361, table III cited in the application --	1-3,6,7,11, 26,27
A	Chemical and Pharmaceutical Bulletin, vol. 35, no. 6, 1987, T. Miyamoto et al.: "Pyridone carboxylic acids as antibacterial agents. VIII. An alternative synthesis of enoxacin via fluoronicotinic acid derivatives", pages 2280-2285 cited in the application	1-3,5-7 ./.
<p>* Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"G" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE		

III. DOCUMENTS CONSIDERED TO BE RELEVANT (C CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	GB, A, 1208279 (ICI LTD) 14 October 1970, see claims 1,2; page 1, lines 48-55 --	1,2,4
Y	Journal of the American Chemical Society, vol. 97, no. 25, 10 December 1975, A.I. Meyers et al.: "Oxazolines. XXII. Nucleophilic aromatic substitu- tion on aryl oxazolines. An efficient approach to unsymmetrically substituted biphenyls and o-alkyl benzoic acids", pages 7383-7385, see page 7383, right- hand column, formula scheme; page 7384, left-hand column, formula scheme --	23
Y	Chemical Abstracts, vol. 84, no. 3, 19 January 1976 (Columbus, Ohio, US), T.N. Gerasimova et al.: "Interaction of pentafluorobenzoic acid and its esters with alkylmagnesium halides", page 425, abstract no. 16902r, & Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk 1975, (5), 100-6, see the abstract --	23
Y	Bull. Chem. Soc. Jpn, vol. 57, 1984, The Chemical Society of Japan, Y. Inukai et al.: "Ortho-disubstituted F-benzenes. V. Intramolecular hetero- atom-facilitated ortho-substitution of F-benzene derivatives with oxazolinyl and oxazinyl groups", pages 225-231, see page 225, scheme 2; page 226, scheme 3 cited in the application --	23
A	US, A, 4469885 (R.A. MUELLER et al.) 4 September 1984, see example 7 --	23
A	Patent Abstracts of Japan, vol. 10, no. 258 (C-370)(2314), 4 September 1986 & JP, A, 6185349 (NIPPON SHOKUBAI KAGAKU KOGYO CO. LTD) 30 April 1986 --	23
A	Patent Abstracts of Japan, vol. 11, no. 167 (C-425)(2614), 28 May 1987 & JP, A, 6245 (NIPPON SHOKUBAI KAGAKU KOGYO LTD) 6 January 1987 -- ./.	23

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

A,P | EP, A, 0287951 (OTSUKA PHARMACEUTICAL CO. 24  
 LTD) 26 October 1988, see page 44,  
 end product of "reference example 21"  
 -----

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This International search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 12... because they relate to subject matter not required to be searched by this Authority, namely:

See PCT-Rule 39.1 (iv): methods for treatment of the human  
 or animal body by surgery or therapy,  
 as well as diagnostic methods.

2. ☐ Claim numbers ..... because they relate to parts of the International application that do not comply with the prescribed requirements to such an extent that no meaningful International search can be carried out, specifically:

3. ☐ Claim numbers ..... because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this International application as follows:

Please refer to Form PCT/ISA/206 dated 30 May 1989.

1. ☒ As all required additional search fees were timely paid by the applicant, this International search report covers all searchable claims of the International application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this International search report covers only those claims of the International application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

# ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 8900278  
SA 26813

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 17/11/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4341784	27-07-82	JP-A- 56049382	02-05-81
		AU-B- 534288	19-01-84
		AU-A- 6257080	09-04-81
		CA-A- 1147731	07-06-83
		EP-A, B 0027752	29-04-81
EP-A- 0207420	07-01-87	JP-A- 62234082	14-10-87
EP-A- 0242789	28-10-87	AU-A- 7190987	29-10-87
		JP-A- 63045261	26-02-88
		ZA-A- 8702874	12-10-87
US-A- 4146719	27-03-79	JP-A- 53141286	08-12-78
		AU-B- 512940	06-11-80
		AU-A- 3271378	02-08-79
		BE-A- 863429	16-05-78
		CA-A- 1178961	04-12-84
		DE-A, B, C 2804097	23-11-78
		FR-A, B 2391210	15-12-78
		GB-A- 1574285	03-09-80
		SE-B- 436129	12-11-84
		SE-A- 7800970	17-11-78
US-A- 4649144	10-03-87	JP-A- 60260577	23-12-85
		JP-A- 60028978	14-02-85
		DE-A- 3470420	19-05-88
		EP-A, B 0132845	13-02-85
US-A- 4571396	18-02-86	AU-B- 566984	05-11-87
		AU-A- 4092085	24-10-85
		EP-A- 0159174	23-10-85
		OA-A- 7995	31-01-87
		JP-A- 60260573	23-12-85
GB-A- 1208279	14-10-70	None	
US-A- 4469885	04-09-84	AU-A- 1122288	05-05-88
		AU-B- 571513	21-04-88
		AU-A- 2778584	15-11-84

EP 0 400 000

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

# ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 8900278  
SA 26813

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 17/11/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 4469885		CA-A, C 1235426	19-04-88
		EP-A, B 0126372	28-11-84
		JP-A- 59210046	28-11-84
		US-A- 4501895	26-02-85
EP-A- 0287951	26-10-88	None	

EP 0 287 951 A1

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82